

Short Communication

Cryptosporidium parvum infection in Calves from an animal farm in Slovakia

Kristína Mravcová¹, Gabriela Štrkolcová^{1*}, Mária Goldová¹, Rastislav Mucha²

¹Institute of Parasitology, University of Veterinary Medicine and Pharmacy in Košice, ²Institute of Neurobiology of Biomedical Research Centre, Slovak Academy of Sciences, Kosice, Slovak Republic

*Corresponding author

Gabriela Štrkolcová, Institute of Parasitology, University of Veterinary Medicine and Pharmacy in Košice, Komenského 73, 041 81 Košice, Slovak Republic, Tel: +421-907-507-473; E-mail: gabriela.strkolcova@uvlf.sk

Submitted: 04 December 2018

Accepted: 03 January 2019

Published: 04 January 2019

ISSN: 2379-948X

Copyright

© 2019 Štrkolcová et al.

OPEN ACCESS

Keywords

- Cryptosporidiosis
- Molecular methods
- Diarrhea
- Calf
- Zoonoses

Abstract

Cryptosporidium spp. is opportunistic pathogen that infect a wide range of animals, including mammals and birds. In the present paper we focused our attention on faecal samples of calves younger than 35 days bred on a cattle farm in Slovakia. The total of collected faecal was 54 and applying the Ziehl-Neelsen acid resistant staining technique, we confirmed the presence of oocysts in 31.5% of smears. And for confirmation was used molecular methods with identification of the *Cryptosporidium parvum* species and on the basis of the gp60 gene we confirmed two zoonotic subtypes, IIdA17G1 and IIdA17G1R1.

INTRODUCTION

Cryptosporidium are intracellular protozoa with a broad host range, cosmopolitan prevalence and zoonotic potential [1]. Until now, 38 valid species of *Cryptosporidium* genus have been described; out of them, more than 20 were reported in humans while *Cryptosporidium hominis* and *Cryptosporidium parvum* are responsible for most human infections [2,3,4]. *Cryptosporidium* spp. represent a significant cause of diarrhoea in farm animals all over the world. The main reservoirs of pathogenic species *Cryptosporidium* spp. include farm animals, especially beef cattle [5]. The impact of a parasite is the strongest in young cattle animals with insufficient colostral immunity. Getting into contact with infected animals represents a risk also for farm staff, veterinary doctors, immune deficient and immune suppressed individuals [6,7,8]. Oocysts are most frequently transmitted directly via the faecal-oral route [9] but they can also be transmitted through contaminated water, feeds, or tools [10,11]. Calves very often become infected within the first week of their lives [age ≤ 30 days] [12]. Typical findings include watery diarrhoea, sometimes accompanied with lethargy, fever, dehydration, and overall poor condition of an animal. Infection may disappear spontaneously within 1-2 weeks; in some cases, it may have a fatal end [13,14]. The species that commonly occur in beef cattle include *Cryptosporidium parvum*, *Cryptosporidium bovis*, *Cryptosporidium ryanae* and *Cryptosporidium andersoni* [15]. The distribution of such species depends on their age: *C. parvum* prevails in pre-weaned calves [up to 2 months of age] [16,17]; *C. bovis* and *C. ryanae* are detected in young cattle animals [3-

12 months] [18,19,20] and *C. andersoni* mostly occurs in adult beef cattle older than 12 months [21]. Pre-weaned calves are therefore regarded as important reservoirs of *C. parvum* oocysts that are infectious for humans [22]. From the epidemiology point of view, in *Cryptosporidium* species it is very important to perform a molecular analysis. Due to the highly polymorphic GP60 gene, numerous subtype families of *C. parvum* have been described. The subtyping is based on the sequence analysis of 60-kDa glycoprotein gene [GP60] containing highly polymorphic microsatellite sequences with variable number of TCA and TCG repeats. The analysis of GP60 gene proved that approximately 98% of *Cryptosporidium* isolates from calves belong to the IId subtype family that is of the zoonotic nature, and it may also be identified in humans [23]. IId and IId subtype families prevail in animals and humans all over the world [24]. These families have been detected in Europe [Hungary, Germany, Portugal, Sweden, Ireland, Spain, Belgium, Romania, United Kingdom, Netherlands, Serbia and Montenegro] [25,26,27], Asia [Kuwait, Iran, Jordan, India, Malaysia and China] [28,29], Egypt, and Australia [30].

Slovakia now belongs to the countries with confirmed presence of IIdA17G1R1 subtype in beef cattle [31] and in oncological patients [32]. Our objective was to identify subtypes of *Cryptosporidium* spp. in calves younger than 35 days, as currently there are very little records on the prevalence of this species in calves in Slovakia. There is only one paper published in the English language, i.e. by Danišová et al., 2016 [5], which confirmed the 70 % prevalence of *C. parvum* in calves younger than 1 month.

The purpose of the present paper was to identify the species and subtypes occurring in calves in Slovakia.

MATERIALS AND METHODS

On a farm in Zemplínska Teplica village, Trebišov District, we collected 54 samples of faeces from calves aged less than 35 days; the samples were collected in the period from October 2017 to November 2018. Faecal smears were stained using Ziehl-Neelsen stain method [33]. The stained oocysts appeared to be spherical formations of pink colour on the blue-green background. The amplification of the SSU rRNA segment sized 820bp was carried out using the nested PCR [34] which confirmed the presence of *C. parvum* species. Subsequently, the subtyping of 5 selected products was carried out. It was based on the sequence analysis of the GP60 locus according to Alves et al. [2003] [35]. DNA extraction was carried out using the commercial ZR Fecal DNA MiniPrep™ kit [Zymo Research, USA] according to the manual recommended by the manufacturer. The reaction mixture for a PCR reaction contained 50 µl of the mixture [Master Mix "OneTaq 2x MM" [New England Biolabs]], primers [forward and reverse] in the concentration of 33µM, H₂O, and DNA. The amplification of the segment sized SSU rRNA [~ 820 – 850 bp] was carried out using the nested PCR. The content of the second PCR reaction [nest 2] was similar to the first one, except for the use of the primary product as a template for the second PCR reaction [nest 2]. The PCR reactions included the positive and negative controls. A microscopically confirmed positive sample was used as a positive control. The program of amplification was as follows: initial denaturation at 94°C for 3 minutes, followed by 35 cycles of 94°C/45 sec, 55°C/45 sec and 72°C/60 sec, followed by a final extension of 72°C/7 min. GP60 gene [850 bp] was also amplified using the nested PCR. Both PCR reactions [nest1, nest2] included 40 cycles consisting of thermal steps of exposure to 95 °C for the period of 45 s, exposure to 52 °C for the period of 45 s, and exposure to 72 °C for 1 minute, with the initial step of denaturation at 95 °C for 3 minutes and the final amplification at 72°C for the period of 10 minutes. Positive 820 – 850 bp PCR products were visualized on a 1.5 % agarose gel, using the Gel red [Biotium Inc CA Hayward USA] and visualized under the UV light.

Secondary PCR products were sent for purification and sequencing to the Microsynth Laboratory in Wien. The final sequences were compared to the known sequences using the BLAST-analysis in the NCBI database [http://blast.ncbi.nlm.nih.gov].

RESULTS AND DISCUSSION

Cryptosporidiosis is a globally prevalent parasitic disease affecting animals and humans [36]. Gastrointestinal tract diseases induced by species of the *Cryptosporidium* genus facilitate transmission among individuals and environmental contamination. A hygiene level significantly affects the ability of excreted oocysts to survive in the external environment where they are infectious for as long as one year [6]. Cryptosporidiosis may develop as an opportunistic infection or a secondary disease. Over the last few years, numerous reports on infections caused by *Cryptosporidium* spp. have been published [2,3,4,5]; this proves that this parasite has spread globally. Cattle may become infected

with four species of *Cryptosporidium* spp.: *C. parvum*, *C. bovis*, *C. andersoni* and *C. Ryanae* [15]. *C. Parvum* is the most important zoonotic species [4]; most frequently occurs in calves aged less than 2 months and is responsible for 80% of *Cryptosporidium* infections in these animals [37]. *C. parvum* belongs to the group of parasites affecting a host's intestine; in beef cattle, it mainly infects jejunum and ileum. Clinical symptoms persist for the period of 4-7 days in form of watery diarrhoea, depression, malnutrition and abdominal pain [38,39]. The prevalence relates to the method and extent of breeding and the density of farm animals population [6].

In Netherlands, the prevalence of *Cryptosporidium parvum* in beef cattle animals younger than three months represents approximately 64% [40]. In the Czech Republic, the prevalence of infected cattle animals younger than 2 months was 21.5% [in 2011]. According to Kaupke, the prevalence of cryptosporidiosis in Poland reached the value of 22.5% in 2015 [41]. In Columbia, the prevalence of infected calves younger than 35 days represented as much as 26.6% [42]. In Slovakia, the infection induced by *C. parvum* was detected in the 27-30% prevalence, often with the 60% mortality [43 – available only in the Slovak language]. Danišová et al., 2016 confirmed the 70 % prevalence of *C. parvum* in calves younger than 1 month [5].

In calves from the farm in Zemplínska Teplica we diagnosed *Cryptosporidium* spp. oocysts in 31.5% of faecal smears [17/54] using the Ziehl Neelsen stain method. Applying the PCR method, we confirmed the zoonotic species *C. parvum*. Five positive faecal samples were used for genotyping. The subtyping of products was carried out on the GP60 locus. The comparison of the obtained sequences and the reference sequences acquired from the GenBank showed that the isolates belonged to IId and IIa subtypes. The IIdA17G1 subtype [GenBank KY499053.1] was identified in four isolates, and the IIaA17G1R1 subtype [GenBank JX258865.1] in one isolate. The IIdA17G1 subtype of *C. parvum* in beef cattle belongs to zoonotic subtypes and in Europe it was also confirmed in humans [44,32], in samples of treated water [44], as well as in small ruminants, for example in Spain [46]. In 2017, Hatalová confirmed the presence of the IIaA17G1R1 subtype in Slovakia in beef cattle [31]. Particularly this subtype that we have identified is associated with zoonotic transmission confirmed in humans e.g. in patient from Great Britain [44]; in 12 patients from Slovakia with oncological diseases [32].

CONCLUSION

Domestic ruminants [especially calves] represent typical hosts for *Cryptosporidium* spp. Within the present study we examined faecal samples collected from calves younger than 35 days. We confirmed the presence of *C. parvum* species and *C. Parvum* zoonotic subtype families IId and IIa. Calves come into close contact with humans [especially with farm workers, veterinary doctors, students of veterinary school...etc] and thus represent a significant risk in terms of cryptosporidiosis spreading. The contaminated water and food by oocysts of *Cryptosporidium* also represents an important pathway of infection.

There is currently no efficient therapy of cryptosporidiosis. It is recommended to initiate symptomatic treatment of diarrhoea and dehydration; in more severe cases, halofuginone is

recommended for affected calves [37]. The most efficient measures include improvement of breeding conditions [clean boxes for calving and calves and separation of infected calves] and regular sanitation of the environment. It is also very important to provide veterinary care and perform regular monitoring of cryptosporidiosis occurrence.

ACKNOWLEDGEMENTS

The study was supported by the grant of the State Agency VEGA No. 1/0536/18.

REFERENCES

- Xiao L, Ryan U. Cryptosporidiosis: an update in molecular epidemiology. *Current Opinion in Infectious Diseases*. 2004; 17: 483-490.
- Ježeková J. *Cryptosporidium testudinissp. n., Cryptosporidium ducimarci* Traversa, 2010 and *Cryptosporidium tortoise* genotype III [Apicomplexa: Cryptosporidiidae] in tortoises. *Folia Parasitologica*. 2016; 63: 1-10.
- Zahedi A. *Cryptosporidium homai* n. sp. [Apicomplexa: Cryptosporidiidae] from the guinea pig [*Caviaporcellus*]. *Veterinary Parasitology*. 2017; 254: 92-101.
- Feng Y, Ryan UM, Xiao L. Genetic Diversity and Population Structure of *Cryptosporidium*. 2018; 34: 997-1011.
- Danišová O, Valenčáková A, Petřincová A. Detection and identification of six *Cryptosporidium* species in livestock in Slovakia by amplification of SSU and GP60 genes with the use of PCR analysis. *Annals of Agricultural and Environmental Medicine*. 2016; 23: 254-258.
- Thompson RCA, Olsen ME, Zhu G, Enomoto S, Abrahamsen MS, Hijjawi NS. *Cryptosporidium* and Cryptosporidiosis. *Advances in Parasitology*. 2005; 59: 77-158.
- Fletcher SM, Stark D, Harkness J, Ellis J. Enteric Protozoa in the Developed World: a Public Health Perspective. *Clin Microbiol Rev*. 2012; 25: 420-449.
- Checkley et al. A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. *Lancet Infect Dis*. 2015; 15: 85-94.
- Adler S, Widerström M, Lindh J, Lilja M. Symptoms and risk factors of *Cryptosporidium hominis* infection in children: data from a large waterborne outbreak in Sweden. *Parasitol Res*. 2017; 116: 2613-2618.
- Food and Agriculture Organization of the United Nations, 2014.
- Thompson RCA, Koh WH, Clode PL. *Cryptosporidium* - What is it? *ScienceDirect*. 2016; 4: 54-61.
- Uga S, Matsuo J, Kono E, Kimura K, Inoue M, Rai SK, Ono K. Prevalence of *Cryptosporidium parvum* infection and pattern of oocyst shedding in calves in Japan. *Vet Parasitol*. 2000; 94: 27-32.
- Tzipori S. Cryptosporidiosis in animals and humans. *Microbiol Rev*. 1983; 47: 84-96.
- Fayer R, Trout JM, Jenkins MC. Infectivity of *Cryptosporidium parvum* Oocysts Stored in Water at Environmental Temperatures. *The Journal of Parasitology*. 1998; 84: 1165-1169.
- Robertson LJ, Björkman C, Axén CH, Fayer R. Cryptosporidiosis in Farmed Animals. *Cryptosporidium: parasite and disease*. Springer, Vienna. 2014; 149-235.
- Brook EJ, Anthony HC, French NP, Christley RM. Molecular epidemiology of *Cryptosporidium* subtypes in cattle in England. *Vet J*. 2009; 179: 378-382.
- Plutzer J, Karanis P. Genotype and subtype analyses of *Cryptosporidium* isolates from cattle in Hungary. *Vet Parasitol*. 2007; 146: 357-362.
- Santin M, Zarlenga DS. A multiplex polymerase chain reaction assay to simultaneously distinguish *Cryptosporidium* species of veterinary and public health concern in cattle. *Vet Parasitol*. 2009; 166: 32-37.
- Fayer R, Santin M, Trout JM. *Cryptosporidium ryanae* n. sp. [Apicomplexa: Cryptosporidiidae] in cattle [*Bostaurus*]. *Vet Parasitol*. 2008; 156: 191-198.
- Muhid A, Robertson I, Ng J, Ryan U. Prevalence of land management factor contributing to *Cryptosporidium* sp. infection in pre-weaned and post-weaned calves in Johor, Malaysia. *Exp Parasitol*. 2011; 127: 534-538.
- Khan SM, Debnath Ch, Pramanik AK, Xiao L, Nozaki T, Ganguly S. Molecular characterization and assessment of zoonotic transmission of *Cryptosporidium* from dairy cattle in West Bengal, India. *Vet Parasitol*. 2010; 171: 41-47.
- Grinberg A, Pomroy WE, Squires RA, Scuffham A. Retrospective cohort study of an outbreak of cryptosporidiosis caused by a rare *Cryptosporidium parvum* subgenotype. *Epidemiology and Infection*. 2011; 139: 1542-1550.
- Ryan U, Fayer R, Xiao L. *Cryptosporidium* species in humans and animals: current understanding and research needs. Cambridge University Press. 2014; 141: 1667-1685.
- Li F. Prevalence and molecular characterization of *Cryptosporidium* spp. and *Giardia duodenalis* in dairy cattle in Beijing, China. *Vet Parasitol*. 2016; 219: 61-65.
- Imre K. Zoonotic *Cryptosporidium parvum* in Romanian newborn lambs [*Ovisaries*]. *Vet. Parasitol*. 2013; 191: 119-122.
- Insulander M, Silverlas C, Lebbad M, Karlsson L, Mattsson JG, Svenungsson B. Molecular epidemiology and clinical manifestation of human cryptosporidiosis in Sweden. *Epidemiol. Infect*. 2013; 141: 1009-1020.
- Vieira PM. Molecular characterization of *Cryptosporidium* [Apicomplexa] in children and cattle in Romania. *Folia Parasitol*. 2015; 62: 1-4.
- Wang R. Characteristic of *Cryptosporidium* transmission in pre-weaned dairy cattle in Henan, China. *J. Clin. Microbiol*. 2011; 49: 1077-1082.
- Iqbal A, Lim YA, Surin J, Sim BL. High diversity of *Cryptosporidium* subgenotypes identified in Malaysian HIV/AIDS individuals targeting gp60 gene. *PLoS One*. 2012.
- Amer S, Zidan S, Feng Y, Adamu H, Li N, Xiao L. Identity and public health potential of *Cryptosporidium* spp. in water buffalo calves in Egypt. *Vet. Parasitol*. 2013; 191: 123-127.
- Hatalová E, Valenčáková A, Kalinová J, Luptáková L, Danišová O, Húska M. Comparison of PCR primers for the identification of *Cryptosporidium* species and genotypes in calves. *Bulgarian Journal of Veterinary Medicine*. 2017; 20: 154-157.
- Hatalová E. The first of animal genotypes of *Cryptosporidium parvum* in immunosuppressed and immunocompetent humans in Slovakia. Wiley online library. 2018; 1-7.
- Henriksen SA, Pohlenz JFL. Staining of Cryptosporidia by a Modified Ziehl-Neelsen Technique. *Acta Veterinaria Scandinavica*. 1981; 22: 594-596.
- Xiao L. Genetic diversity within *Cryptosporidium parvum* and related *Cryptosporidium* species. *Appl Environ Microbiol*. 1999; 65: 3386-3391.

35. Alves M, Xiao L, Sulaiman I, Lal AA, Matos O, Antunes F. Subgenotype Analysis of *Cryptosporidium* Isolates from Humans, Cattle, and Zoo Ruminants in Portugal. *Journal of Clinical Microbiology*. 2003; 41: 2744-2747.
36. Cacciò SM, Chalmers RM. Human cryptosporidiosis in Europe. *Clinical Microbiology and Infection*. 2016; 22: 471-480.
37. Deplazes P, Eckert J, Mathis A, Samson-Himmelstjerna G, Zahner H. *Parasitology in Veterinary Medicine*. Wageningen Academic Publishers The Netherlands. 2016.
38. Tzipori S, Smith M, Halpin C, Angus KW, Sherwood D, Campbell I. Experimental cryptosporidiosis in calves: clinical manifestations and pathological findings. *Vet Rec*. 1983; 112: 116-120.
39. Fayer R, Gasbarre L, Pasquali P, Canals A, Almeria S, Zarlenga D. *Cryptosporidium parvum* infection in bovine neonates: dynamic clinical parasitic and immunologic patterns. *Int J Parasitol*. 1998; 28:49-56.
40. Toledo RdS, Martins FDC, Ferreira FP, de Almeida JC et al. *Cryptosporidium* spp. and *Giardia* spp. in feces and water and the associated exposure factors on dairy farms. *PLOS*. 2017; 12: 1-20.
41. Kaupke A, Rzezutka A. Emergence of novel subtypes of *Cryptosporidium parvum* in calves in Poland. *Parasitology Research*. 2015; 114: 4709-4716.
42. Avendaño C, Ramo A, Vergara-Castiblanco C, Sánchez-Acedo C, Quílez J. Genetic uniqueness of *Cryptosporidium parvum* from dairy calves in Colombia. *Parasitology Research*. 2018; 117: 1317-1323.
43. Mudroň P, Mudroňová D, Goldová M, Nagy O. *Cryptosporidium parvum* - Unusual cause of chronic diarrhea in dairy cows. *Nezvyklá příčina chronickej hnačky u dojníc*. *InfoVet*. 2003; 3: 33-34.
44. Chalmers RM, Smith RP, Hadfield SJ, Elwin K, Giles M. Zoonotic linkage and variation in *Cryptosporidium parvum* from patients in the United Kingdom. *Parasitol. Res*. 2011; 108: 1321-1325.
45. Lobo ML, Xiao L, Antunes F, Matos O. Occurrence of *Cryptosporidium* and *Giardia* genotypes and subtypes in raw and treated water in Portugal. *Lett. Appl. Microbiol*. 2009; 48: 732-737.
46. Díaz P. *Cryptosporidium* species and subtype analysis in diarrheic pre-weaned lambs and goat kids from north-western Spain. *Parasitol. Res*. 2015; 114: 4099-4105.

Cite this article

Mravcová K, Štrkolcová G, Goldová M, Mucha R (2019) *Cryptosporidium parvum* infection in Calves from an animal farm in Slovakia. *J Vet Med Res* 6(1): 1169.