

## Mini Review

# Paget's Disease of the Vulva and Penis: A Mini Review

John L. Powell\*

*Department of Obstetrics and Gynecology, University of North Carolina School of Medicine, USA***\*Corresponding author**

John L. Powell, Department of Obstetrics and Gynecology, University of North Carolina School of Medicine, 10 Crane Street, Wrightsville Beach, N.C. 28480, Tel: 910-256-6812; Email: john.powellatwb@att.net

**Submitted:** 30 October 2018**Accepted:** 06 December 2018**Published:** 07 December 2018**ISSN:** 2475-9430**Copyright**

© 2018 Powell

OPEN ACCESS

**Abstract**

Paget's disease of the vulva appears as a patchy, reddish and whitish, velvety, and eczematous skin lesion. Women with Paget's disease of the vulva are usually white and postmenopausal and complain of localized itching and burning. It is confined to the epithelium in most cases, but invasive disease is present in 13% to 25% of cases. If the disease appears to be intraepithelial, then wide local excision is recommended with a margin of 1 to 2 cm. beyond the clinically visible lesion. The use of intraoperative frozen sections has been reported to reduce the incidence of positive margins on permanent sections and lower the risk of recurrences.

**REVIEW LITERATURE**

Paget's disease is classified, according to location, as mammary or extramammary disease. The original lesion was described by James Paget (1814-1889) in 1874, involved the nipple and areola of the breast in 13 women, who developed underlying ductal adenocarcinoma within three years of presentation [1]. Extramammary Paget's disease most commonly involves the vulva and anus and was first described in 1901, by William Auguste Dubreuilh (1857-1935), professor of dermatology at the University of Bordeaux [2]. Paget's disease appears as a patchy, reddish and whitish, velvety, and eczematous skin lesion. Women with Paget's disease of the vulva are usually white and postmenopausal and complain of localized itching and burning.

Paget's disease of the vulva is of apocrine origin and is confined to the epithelium in most cases. However, invasive disease is present in 13 to 25% of cases, either, as a result of direct invasion through the basement membrane or, less commonly, because of the presence of an underlying apocrine gland adenocarcinoma. Histologically, intraepithelial Paget's disease appears as large, pale cells, often in nests at the tips of the rete ridges. The cells are often seen infiltrating upward in the epithelium, which is hyperkeratotic. The Paget's cells can be located within any of the skin adnexa.

Vulvar Paget's disease occurs with other malignancies in about 35% of patients; the most common of these is breast carcinoma. Other commonly associated cancers are basal cell, rectal, and genitourinary carcinomas. When Paget's disease involves the anus, there is a very high incidence of coexisting rectal cancer. Part of the preoperative work-up should be directed at screening for these cancers.

When a patient is diagnosed with Paget's disease of the

vulva, the area of involvement should be carefully inspected and palpated to detect areas suspicious for invasive cancer. If the disease clinically appears to be intraepithelial, then a wide local excision is performed, including a small amount of subcutaneous tissue. A well-known characteristic of intraepithelial Paget's disease is histologic extension far beyond that which is clinically apparent. Some authors advocate resection up to 1 to 2 cm beyond the clinically visible lesion. However, they report recurrence rates of 29.6% to 42% [3,4]. They also report up to a 52% incidence of positive margins on permanent sections [4]. Some authors advocate the use of intra-operative frozen sections to define the extent of surgery necessary in order to reduce the incidence of positive margins and recurrences to as low as 11% [5,6]. Frozen sections will not guarantee a 0% recurrence rate, but can significantly reduce them. Intraoperative mapping involves excising strips of tissue along the initially planned margins and sending them for frozen section. While waiting for the frozen section reports, the central lesion is excised. Other methods that have been reported to be useful in ensuring clear margins include Mons micrographic surgery [7] and fluorescein dye with ultraviolet light [8]. Colposcopy and toluidine blue staining are not helpful [9]. An experimental technique reported to be potentially useful in evaluating histologically negative margins is the application of a panel of monoclonal antibodies that may detect occult Paget cells [10].

When intraepithelial Paget's disease extends far beyond that which is clinically apparent, very extensive excision is necessary to obtain clear margins.<sup>6</sup> Primary closure of vulvar wounds is desirable, but may not be possible in such cases. Possible means of dealing with such cases include skin grafting, laser vaporization, and pedicle flap reconstruction [6,11,12]. Paget's disease can recur in a skin graft. Topical 5-fluorouracil or bleomycin, administered either pre- or postoperatively to treat

recurrent disease [13]. Topical imiquimod 5% cream has also been used [14]. Management of recurrent intraepithelial Paget's disease of the vulva is similar to that of the primary lesion, with excision of at least the clinically evident disease. When vulvar Paget's disease is found to contain an invasive component, then the treatment is radical surgery [6]. Radiotherapy has been used as an alternative therapeutic approach for patients with extensive inoperable disease or medical contraindications for the surgical approach [15-18]. Salvage radiotherapy can be given for regional lymph node metastases, inoperable loco-regional recurrence, and in those who refuse additional surgery.

Extramammary Paget's disease affecting the scrotum and penis was first described by Henry Radcliffe Crocker (1845-1909) in 1888 [19]. Crocker was a dermatologist at the University College Hospital and Middlesex School of Medicine in London. The mainstay of treatment is wide surgical excision [20,21]. In a series of 36 patients, 13 (36.1%) underwent local wide excision with frozen section evaluation of the margins and only 1 patient (7.7%) had a positive surgical margin on permanent sections [21]. However, 23 (63.9%) underwent local wide excision with margins of 1 to 2 cm only by gross examination, but 17 (73.9%) of them had positive margins on permanent sections. Topical 5-fluorouracil has been used for intra-epithelial disease [22] and combined chemo radiotherapy has been successfully used for regionally advanced unresectable disease [23].

## REFERENCES

- Paget J. On disease of the mammary areola preceding cancer of the mammary gland. *St Barth Hosp Rep* 1874; 10: 87-89.
- Dubreuilh W. Paget's disease of the vulva. *Br J Dermatol*. 1901; 13: 407-413.
- Sopracordevole F, Di Giuseppe J, De Piero G, Canzonieri V, Buttignol M, Giorda G, et al. Treatment of Paget disease of the vulva: Prognostic significance of stromal invasion and surgical margin status. *J Low Gen Tract Dis*. 2016; 20: 184-188.
- Pierie JP, Choudry U, Muzikansky A, Finkelstein DM, Ott MJ. Prognosis and management of extramammary Paget's disease and the association with secondary malignancies. *J Am Coll Surg*. 2003; 196: 45-50.
- Stacy D, Burrell MO, Franklin EW. Extramammary Paget's disease of the vulva and anus: Use of intra-operative frozen section margins. *Am J Obstet Gynecol* 1986; 155: 519-523.
- Powell JL. Paget Disease of the Vulva: Significance of Surgical Margin Status. *J Low Genit Tract Dis*. 2016; 20: e53.
- Coldiron BM, Goldsmith BA, Robinson JK. Surgical treatment of extramammary Paget's disease: A report of six cases and a reexamination of Mohs micrographic surgery compared with conventional surgical excision. *Cancer*. 1991; 67: 933-938.
- Misas JE, Cold CJ, Hall FW. Vulvar Paget's disease: Fluorescein-aided visualization of margins. *Obstet Gynecol*. 1991; 77: 156-159.
- Friedrich EG Jr, Wilkinson EJ, Steingraeber PH, Lewis JD. Paget's disease of the vulva and carcinoma of the breast. *Obstet Gynecol*. 1975; 46: 130-134.
- Ganji P, Giraldo KA, Lampe B, Nadji M. Vulvar Paget's disease. Is immunocytochemistry helpful in assessing the surgical margins? *J Reprod Med*. 1990; 35: 1002-1004.
- Ewing TL. Paget's disease of the vulva treated by combined surgery and laser. *Gynecol Oncol*. 1991; 43: 137-140.
- Valentine BH, Arena B, Green E. Laser ablation of recurrent Paget's disease of vulva and perineum. *J Gynecol Surg*. 1992; 8: 21-24.
- Waring WG, Roberts JA, Lagasse LD, Berman ML, Ballon SC, Moore JG, et al. Treatment of recurrent Paget's disease of the vulva with topical bleomycin. *Cancer*. 1978; 41: 10-11.
- Feldmeyer L, Kerl K, Kamarashev J, de Viragh P, French LE. Treatment of vulvar Paget disease with topical imiquimod: a case report and review of the literature. *J Dermatol Case Rep*. 2011; 5: 42-46.
- Son SH, Lee JS, Kim YS, Ryu MR, Chung SM, Namkoong SE, et al. The role of radiation therapy for the extramammary paget's disease of the vulva ; experience of 3 cases. *Cancer Res Treat*. 2005; 37: 365-369.
- Tolia M, Tsoukalas N, Sofoudis C, Giaginis C, Spyropoulou D, Kardamakis D, et al. Primary extramammary invasive Paget's vulvar disease: What is the standard, what are the challenges and what is the future for radiotherapy? *BMC Cancer*. 2016; 16: 563.
- Hata M, Koike I, Wada H, Miyagi E, Odagiri K, Minagawa Y, et al. Definitive radiation therapy for extramammary Paget's disease. *Anticancer Res*. 2012; 32: 3315-3320.
- Karam A, Dorigo O. Treatment outcomes in a large cohort of patients with invasive extramammary Paget's disease. *Gynecol Oncol*. 2012; 125: 346-351.
- Crocker HR. Paget's disease affecting the scrotum and penis. *Trans Pathol Soc London*. 1889; 40: 187-191.
- Ng LG, Yip SK, Tan PH. Extramammary Paget's disease of the scrotum. *Urology*. 2001; 58: 105.
- Yang WJ, Kim DS, Im YJ, Cho KS, Rha KH, Cho NH, et al. Extramammary Paget's disease of penis and scrotum. *Urology*. 2005; 65: 972-975.
- Bewley AP, Bracka A, Staughton RC, Bunker CB. Extramammary Paget's disease of the scrotum: treatment with topical 5-fluorouracil and plastic surgery. *Br J Dermatol*. 1994; 131: 445-446.
- Voigt H, Bassermann R, Nathrath W. Cytoreductive combination chemotherapy for regionally advanced unresectable extramammary Paget carcinoma. *Cancer*. 1992; 70: 704-708.

### Cite this article

Powell JL (2018) Paget's Disease of the Vulva and Penis: A Mini Review. *Ann Clin Cytol Pathol* 4(8): 1126.