

Research Article

Sudden Death of Cattle Caused By *Babesia Bovis* Transmitted By *Rhipicephalus Microplus* (Acari: Ixodidae) Ticks

Jin Luo^{1*}, Zhen Chen², Kai Fei Guo², Deng Ji Lou², Rong Han⁴, Li Yan Che^{1,3}, Meng Yang Gou¹, Gui Quan Guan¹, Jian Xun Luo¹, Hong Yin¹, and Guang Yuan Liu¹

¹State Key Laboratory of Veterinary Etiological Biology, Key Laboratory of Veterinary Parasitology of Gansu Province, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Science, China

²School of Chemistry, Biology and Environment, Yuxi Normal University, China

³College of Animal Sciences, Tarim University / Key Laboratory of Tarim Animal Husbandry Sciences and Technology, Xinjiang Production and Construction Corps, China

⁴Central Agricultural Radio and Television School, China

***Corresponding author**

Jin Luo, State Key Laboratory of Veterinary Etiological Biology, Key Laboratory of Veterinary Parasitology of Gansu Province, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Science, China

Submitted: 11 August, 2023

Accepted: 26 August, 2023

Published: 28 August, 2023

ISSN: 2333-6684

Copyright

© 2023 Luo J, et al.

OPEN ACCESS**Keywords**

- Ticks
- *Rhipicephalus Microplus*
- *Babesia Bovis*
- Acute Death

Abstract

In this study, a case of acute cattle death was investigated. The pathogen was identified, and its molecular characteristics and vector were analysed. This study provides a reference for the prevention and control of babesiosis and the healthy breeding of cattle. In this study, unengorged and engorged *Rhipicephalus microplus* ticks were collected from the Chongqing area. The unengorged ticks were cultured on cattle under laboratory conditions, and the engorged ticks were cultured to lay eggs. In the process, the cattle suddenly died at 12 days from the bite of an unengorged *R. microplus* tick. In addition, the larvae hatched from *R. microplus* eggs, which were cultured on the other cattle, and the experimental cattle died in approximately 14 days. Blood was collected from a second dying and stored at 4°C for one week. Two millilitres of anticoagulated blood was injected subcutaneously into the third cow without piriformis infection. On the fourth day, the body temperature rose to 41°C with slight lymphadenopathy. On the fifth day, the cow suddenly fell and died approximately 4 hours later. DNA was extracted from the blood of all dead cattle and amplified by PCR with piriformis universal primers. The results showed that the cattle were infected with *Babesia bovis*. The phylogenetic tree based on 18S rRNA showed that the Chongqing strain of *B. bovis*, which caused the death of cattle, was closely related to the Yunnan strain in China and had the same taxonomic status as the Spanish strain. This case report will draw people's attention to *B. bovis* once again, and positive control measures should be taken to reduce the losses of farmers to achieve the goal of healthy breeding.

INTRODUCTION

Babesia bovis is a tick-borne intracellular protozoan parasite that causes the most pathogenic form of bovine babesiosis [1]. It can lead to high fever, mental depression, loss of appetite, and even death, causing serious harm to the world cattle industry and human public health [2]. Many researchers have carried out a series of prevention and control studies based on *B. bovis*. The use of chemicals and acaricides has caused serious environmental pollution and resistance to pathogens, and they can no longer adapt to current social development and health needs [3]. Therefore, vaccines have become the focus of research. At present, live attenuated vaccines have achieved good efficiency in disease prevention and animal immune protection and have been widely used in Australia, Argentina, Brazil, Uruguay and Israel [4]. In addition, the technology of *in vitro* culture of *B. bovis* is convenient for biochemical, immunological and antigen source research [5]. In several studies conducted in Mexico, an attenuated

vaccine of *B. bovis in vitro* produced good protection for host animals, but the disadvantage of this method is that it requires a large-scale culture of *B. bovis*. In recent years, the screening of immune molecules has also become an important means to control *B. bovis*. Examples include natural killer cells from the spleen, antigens identified by immune CD4+ T lymphocytes, MHC class II restricting elements, etc. [6]. Ideally, the *B. bovis* vaccine must induce a humoral immune response, which is characterized by neutralizing antibodies and cellular Th1 immune responses against conservative epitopes. The AMA-1, MSA-2c and RAP-1 proteins have been studied in *B. bovis*, and antibodies to these proteins have shown good neutralizing effects, suggesting the role of B cells and T cell epitopes in the immune response. However, whether there are conserved peptides in the epitopes of B cells and T cells in all strains and their role in producing lasting immunity remain to be determined [7]. The spread and epidemic characteristics of *B. bovis* are important components in the prevention and control of bovine worm disease caused by

this pathogen. Studies have shown that *R. microplus* ticks are an important biological medium for *B. bovis* [8,9]. Therefore, based on the prevalence of ticks, the prevalence of *B. bovis* is seasonal and time-sensitive.

In this study, *R. microplus* were collected from Chongqing city, China, and cultured in the laboratory on cattle. However, the cattle died after being bitten by the ticks. Here, the blood of infected cattle was collected, the pathogens were detected with piriformis universal primers for PCR, and positive PCR products were used for sequencing analysis. Based on the sequencing results, it is speculated that the pathogen may have caused the cows' deaths. To further clarify the prevalence of this pathogen in China, ticks in the main provinces experiencing an epidemic of *R. microplus* were investigated, and the status of their infection with the pathogen was analysed to provide a reference for the formulation of prevention and control measures and early warning of the pathogen.

MATERIALS AND METHODS

Ethics Statement

All animal experiments were performed according to the protocols approved by the Animal Care and Use Committee of the Lanzhou Veterinary Research Institute (permit number 2023-03).

Tick Collection and DNA Extraction

Engorged adult *R. microplus* ticks were collected from cattle in Chongqing city and identified by morphology in the Department of Veterinary Parasitology, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences. These ticks were maintained at a temperature of $30 \pm 2^\circ\text{C}$ and relative humidity of $80 \pm 5\%$. Some of the ticks were immediately placed into Phosphate Buffered Saline (PBS) and washed twice in a solution containing 0.133 M NaCl, 1.11% Sodium Dodecyl Sulfate (SDS) and 0.0088 M Ethylenediaminetetraacetic Acid (EDTA). These ticks were mixed and stored in liquid nitrogen until ground using a mortar and pestle with liquid nitrogen, and genomic DNA (gDNA) was extracted with a QIAamp DNA Mini Kit (QIAGEN, China) following the manufacturer's instructions. Other unengorged ticks were used to bite experimental cattle.

Cattle Experiment

Experimental cattle had their back hair removed and cloth bags with openings at both ends that had been sewn in advance were glued to the sheared areas. The above mentioned hatched *R. microplus* were released into the cloth bag to bite the cattle to further verify the nonaccidental case of cattle death.

Blood Genome Extraction and PCR Amplification

The gDNA was extracted from cattle blood with a Blood & Tissue Kit (Lot 163043916) from QIAGEN, China, following the manufacturer's instructions. The gDNA of cow blood and ticks was analysed using common primers of piroplasma, 989/990

(Theileria) and Prio-As/S (Babesia) primers [10,11]. The reaction system was 50 μL volume: 36.5 μL of sterilized deionized water, 4 μL of dNTP mixture (2.5 mmol/L), 5 μL of $10\times$ PCR buffer, 1 μL of each primer (25 pmol/ μL), 2 μL of gDNA template, and 0.5 μL of rTaq DNA polymerase (5 U/ μL). The reaction conditions were as follows: 94°C denaturation for 4 min; 40 cycles of 94°C for 1 min, 57°C for 1 min, and 72°C for 1 min; and extension at 72°C for 7 min. The products were electrophoresed in agarose gel at 10 g/L. The PCR products were purified using a TaKaRa Agarose Gel DNA Purification Kit Ver. 2.0 (TaKaRa, Dalian, China), and the amplified products were ligated into the vector pMD[®]19-T (TaKaRa, Dalian, China). The positive clones were sequenced with vector-specific primers (T7 and SP6) by Sangon (Shanghai, China).

Sequence Analysis and Phylogenetic Tree

For homology analyses, the 18S rRNA sequences were searched against the BlastN/X (NCBI) (<http://www.ncbi.nlm.nih.gov/BLAST/>). These sequences from ticks and cattle blood were aligned with previously identified 18S rRNA sequences. Nucleotide sequences from other species retrieved from NCBI GenBank were aligned using Clustal version 1.81. A Neighbour-Joining (NJ) phylogenetic tree was constructed using Molecular Evolutionary Genetics Analysis (MEGA) version 11.0 [12], and the likelihood of branching was tested using bootstrap resampling (1000 pseudoreplicates).

Sample Testing

A total of 1768 wild ticks from Guangxi, Yunnan, Qinghai, Fujian and Gansu provinces were analysed using specific primers for *B. bovis* by PCR [13].

RESULT

Pathological Replication

In the experiment, unengorged *R. microplus* ticks from Chongqing were released to healthy cattle not infected by piroplasma. Other engorged ticks were cultured under laboratory conditions to lay eggs, and the hatched larvae were released to healthy cattle not infected by piroplasma. As a result, all the cattle died suddenly in approximately two weeks. In the process, the ticks and cattle received attention; unfortunately, the experimental cattle died suddenly after approximately 13 days, and they showed some clinical symptoms, such as high fever, enlarged anterior shoulder lymph nodes, mental fatigue and loss of appetite. Blood was collected from ill cattle, and 7% Dimethyl Sulfoxide (DMSO) was added to the blood for storage in liquid nitrogen for later use. Similarly, the blood was inoculated into other experimental cattle, which also exhibited the same clinical symptoms as the previous cattle and ultimately died.

Pathological Examination

An autopsy of sick and dead cows showed thin blood and poor coagulation; tissue oedema, yellow stains on the organs; flushing of the intestines and wrinkling of gastric mucosa with point bleeding. There were bleeding points in the inner and external

membranes of the heart, and the myocardial texture was soft; the liver and spleen were yellow-brown and obviously swollen; the kidneys were light red and yellow and enlarged, and there was dot bleeding on their surface (Figure 1).

PCR Detection and Sequencing

Specific primers for *B. bovis* were used to amplify 18S rRNA by PCR from the blood of dead cattle and *R. microplus* ticks. The results of agarose gel electrophoresis and sequencing showed that the pathogen that caused the death of the cattle at this time was *B. bovis* transmitted by *R. microplus*, and the blood smear also showed the typical morphology of *B. bovis* (Figure 2).




Figure 1 Pathological features of the anatomical tissues of cattle with *Babesia bovis*.

- A. There is a small amount of nasal fluid in the nasal cavity, mixed with mucus and blood;
- B. The mucosa of the intestine and wrinkled stomach is flushed with spot-like bleeding;
- C. The blood is thin and poorly clotting;
- D. The spleen is markedly enlarged with petechial bleeding on the surface.




Figure 2 Morphological characteristics of *Babesia bovis*. Note: The location indicated by the arrow is the morphology of *Babesia bovis* infected bovine red blood cells.

Sequence Analysis and Evolutionary Tree

The 18S rRNA gene sequence of *B. bovis* was obtained and compared with the 18S rRNA nucleic acid sequence of different local strains of this species in the GenBank database. The results show that these sequences are highly conserved, and their identity is more than 92%. The similarity comparison results are shown in (Table 1).

MEGA 11.0 software was used to evaluate evolutionary relationships of the *B. bovis* 18S rRNA gene and showed that the Chongqing strain of *B. bovis* in this case was similar to the Yunnan strain, China (KY805830), and they have the same taxonomic status as the Spanish strain (FJ426364) (Figure 3). On the other hand, the 18S rRNA of *B. bovis* can be divided into two main genotypes, both of which have been found in China.

Sample Detection

In this study, a total of 1768 *R. microplus* were collected in Guangxi, Fujian, Qinghai, Gansu, Chongqing, Shandong, Henan and Yunnan provinces, and these samples were tested for *B. bovis* infection. The results show that the population density of *R. microplus* in Gansu and Yunnan is relatively high, at 358/per square kilometre in Gansu and 356/per square kilometre in Yunnan; the population density of *R. microplus* in Henan is relatively low, with an average of 83/per square kilometre. In these data, the infection rate was high in Chongqing at 21.35%, 18.44% in Gansu Province, and 17.70% in Yunnan Province (Table 2).

DISCUSSION

B. bovis is the most economically important tick-borne disease and is mainly transmitted by *R. microplus* ticks [14]. The disease causes substantial economic losses through high animal mortality, low growth rates, reduced milk production




Figure 3 Neighbour-joining tree showing the phylogenetic relationships of the 18S rRNA based on nucleotide sequences of *Babesia bovis*.

Note: The numbers represent the percentage of 1000 replicates (bootstrap support) for which the same branching patterns were obtained.

Table 1: Information on the 18S rRNA gene sequences of different *Babesia bovis* strains from NCBI.

Species	Isolate	Accession No.	Identity	Species	Isolate	Accession No.	Identity
<i>B. bovis</i>	Chongqing	No		<i>B. bovis</i>	Yunnan	KY805830	96.39
<i>B. bovis</i>	Yunnan	MN252438	96.31	<i>B. bovis</i>	Hunan	JQ723013	95.16
<i>B. bovis</i>	Yunnan	MN252440	95.50	<i>B. bovis</i>	North	KP710223	94.48
<i>B. bovis</i>	Shannxian	AY603398	95.41	<i>B. bovis</i>	Turkey	KP745628	94.11
<i>B. bovis</i>	Jilin	MN900523	95.82	<i>B. bovis</i>	Argentina	MH569534	94.49
<i>B. bovis</i>	Spain	FJ426364	94.84	<i>B. bovis</i>	Germany	EF458213	94.49
<i>B. bovis</i>	South Africa	MH257724	96.23	<i>B. bovis</i>	Australia	JQ437260	95.08
<i>B. bovis</i>	USA	MH046907	96.15	<i>B. bovis</i>	Cuba	MN053043	95.90
<i>B. bovis</i>	Portugal	KP745628	94.11	<i>B. bovis</i>	Bolivia	LC645224	95.54
<i>B. bovis</i>	India	KF928959	95.82	<i>B. bigemina</i>	Yunnan	MH208614	

Table 2: *Babesia bovis* was detected from *Rhipicephalus microplus* ticks in China by PCR.

Province	<i>Rhipicephalus Microplus</i> ticks (Number)	<i>Babesia Bovis</i> Positive (%)
Guangxi	186	12.40 (n = 23)
Yunnan	356	17.70 (n = 63)
Fujian	136	0
Qinghai	225	10.67 (n = 24)
Gansu	358	18.44 (n = 66)
Chongqing	178	21.35 (n = 38)
Shandong	246	9.75 (n = 24)
Henan	83	0
Total	1768	13.46 (n = 238)

in diseased or rehabilitated animals, and the direct costs of tick control and disease treatment. In severe cases, it can lead to the death of an infected animal [15]. In this study, a case of acute death of multiple cattle by bites from field-captured *R. microplus* was discussed. Blood smears and PCR were used to detect pathogens. To observe whether the death of cattle is caused by blood protozoa, we inoculated the blood of previously dying cattle into another cow. However, for approximately 6 days, the cow experienced elevated body temperature, loss of appetite and mental depression. On the 8th day, the cow fell to the ground and could not stand. It died in approximately 4 hours, leaving dark red blood from the nasal cavity. The anatomy showed that there was no obvious change in the lungs, but the spleen was swollen, and the liver also had a small number of bleeding points. According to the clinical features of cattle disease, as well as blood smears, PCR sequencing studies showed that the death was caused by *B. bovis* infection. Pathological anatomy also showed the same pathological features as *B. bovis* disease, such as the gastric surface mucosa being flushed with punctate haemorrhage, poor blood coagulation, serious spleen, and haemorrhage. These pathological changes are the main clinical indicators of Babesia infection [16]. These lesions may also be the main cause of death.

The occurrence and development of *B. bovis* are closely related to the distribution and prevalence of ticks. Available studies have shown that *R. microplus* is mainly distributed in Asia [17], India [18], West Africa [19], and South America [20]. This also implies that the transmission of the *R. microplus* carrying the *B. bovis* pathogen is linked with animal migration and the long-distance transmission of the livestock and poultry trade. This results in the presence of the same strain or the same genotype

in different regions. It also provides a basis for analysing the migration relationship and evolutionary characteristics of different local strains and species. Here, the 18S rRNA of *B. bovis* was amplified in *R. microplus* from Chongqing city, and the 18S rRNA sequences were used to construct the phylogenetic tree. The results showed that the Chongqing strain had high identity with the Yunnan strain (KY805830) in China, but they had the same taxonomic status as the Spanish strain (FJ426364), and the Spanish strain had an earlier evolutionary history. This also suggests that the pathogenic strains will be prevalent in different regions along with animal migration or livestock trade. On the other hand, *B. bovis* 18S rRNA can be divided into two major genotypes (I and II), a result that is consistent with the findings of Ramos CM (2010) [21]. Type I is mainly distributed in China, Spain, West Africa, South America and Australia, and type II is relatively limited, mainly distributed in Turkey, China and Germany. Although the genotype of the *B. bovis* species is simple, its distribution is extremely widespread. The frequent exchanges of livestock and poultry trade around the world have also led to the widespread prevalence of babesiosis, causing certain losses to the local farming industry and threats to human public health security.

The distribution of ticks is closely related to tick-borne diseases. The diseases caused by tick-borne pathogens will occur and develop regularly with tick activity, which shows typical seasonal and spatiotemporal specificity. To understand the distribution of *R. microplus* in different areas in China, *R. microplus* ticks were collected from Guangxi, Fujian, Shaanxi, Gansu, Chongqing, Shandong, Henan, and Yunnan, and *B. bovis* was detected in 1768 ticks using Prio-As/S primers. The results showed that the population density of *R. microplus* varied greatly in different regions, with a relatively high density in Gansu and Yunnan and a relatively low density in Henan. Moreover, the infection rate of *B. bovis* transmitted by *R. microplus* also showed a positive correlation with changes in tick population density. Of course, this inference also requires a comprehensive and detailed investigation of the distribution of local *R. microplus* and the epidemic characteristics of *B. bovis* carried by ticks. Therefore, the data on the population density and infection status of *B. bovis* in different regions of China may not be accurate. However, these data have certain reference value for the prevention and control of *R. microplus* and bovine babesiosis and are important for early warning of bovine babesiosis.

CONCLUSION

In conclusion, *B. bovis* has already occurred in vast areas of our country, seriously endangering the healthy development of the local cattle industry. Therefore, the prevention and control of ticks, the reduction of tick breeding population density, and the avoidance of pathogen transmission during animal migration and livestock trade are important measures to prevent the occurrence of *B. bovis*. Healthy animal breeding and scientific management are also important strategies for the prevention and control of bovine babesiosis.

REFERENCES

- Hakimi H, Asada M, Ishizaki T, Kawazu S. Isolation of viable Babesia bovis merozoites to study parasite invasion. *Sci Rep.* 2021; 11(1): 16959. doi: 10.1038/s41598-021-96365-w. PMID: 34417510; PMCID: PMC8379152.
- Giglioti R, Okino CH, Azevedo BT, Wedy BCR, Gutmanis G, Veríssimo CJ, et al. Semi-quantitative evaluation of Babesia bovis and B. bigemina infection levels estimated by HRM analysis. *Ticks Tick Borne Dis.* 2021; 12(5): 101753. doi: 10.1016/j.ttbdis.2021.101753. Epub 2021 Jun 4. PMID: 34134061.
- Cuy-Chaparro L, Reyes C, Díaz-Guiot EV, Moreno-Pérez DA, Patarroyo MA. Developing Anti-Babesia bovis Blood Stage Vaccines: A New Perspective Regarding Synthetic Vaccines. *Int J Mol Sci.* 2023; 24(6): 5219. doi: 10.3390/ijms24065219. PMID: 36982294; PMCID: PMC10049154.
- Alvarez JA, Rojas C, Figueroa JV. An Overview of Current Knowledge on in vitro Babesia Cultivation for Production of Live Attenuated Vaccines for Bovine Babesiosis in Mexico. *Front Vet Sci.* 2020; 7: 364. doi: 10.3389/fvets.2020.00364. PMID: 32671114; PMCID: PMC7332553.
- Álvarez Martínez JA, Figueroa Millán JV, Ueti MW, Rojas-Martínez C. Establishment of Babesia bovis In Vitro Culture Using Medium Free of Animal Products. *Pathogens.* 2021; 10(6): 770. doi: 10.3390/pathogens10060770. PMID: 34205286; PMCID: PMC8235554.
- Brown WC, Norimine J, Knowles DP, Goff WL. Immune control of Babesia bovis infection. *Vet Parasitol.* 2006; 138(1-2): 75-87. doi: 10.1016/j.vetpar.2006.01.041. Epub 2006 Feb 28. PMID: 16510249.
- Hidalgo-Ruiz M, Mejía-López S, Pérez-Serrano RM, Zaldívar-Lelo de Larrea G, Ganzinelli S, Florin-Christensen M, et al. Babesia bovis AMA-1, MSA-2c and RAP-1 contain conserved B and T-cell epitopes, which generate neutralizing antibodies and a long-lasting Th1 immune response in vaccinated cattle. *Vaccine.* 2022; 40(8): 1108-1115. doi: 10.1016/j.vaccine.2022.01.023. Epub 2022 Jan 22. PMID: 35078663.
- Githaka NW, Bishop RP, Šlapeta J, Emery D, Nguu EK, Kanduma EG. Molecular survey of Babesia parasites in Kenya: first detailed report on occurrence of Babesia bovis in cattle. *Parasit Vectors.* 2022; 15(1): 161. doi: 10.1186/s13071-022-05279-7. PMID: 35526030; PMCID: PMC9077973.
- Guglielmo AA. Epidemiology of babesiosis and anaplasmosis in South and Central America. *Vet Parasitol.* 1995; 57(1-3): 109-119. doi: 10.1016/0304-4017(94)03115-d. PMID: 7597777.
- Li Y, Guan G, Ma M, Liu J, Ren Q, Luo J, et al. Theileria ovis discovered in China. *Exp Parasitol.* 2011; 127(1): 304-307. doi: 10.1016/j.exppara.2010.07.002. Epub 2010 Jul 7. PMID: 20619262.
- Gou H, Guan G, Liu A, Ma M, Xu Z, Liu Z, et al. A DNA barcode for Piroplasma. *Acta Trop.* 2012; 124(1): 92-97. doi: 10.1016/j.actatropica.2012.07.001. Epub 2012 Jul 14. PMID: 22796448.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol.* 2018; 35(6): 1547-1549. doi: 10.1093/molbev/msy096. PMID: 29722887; PMCID: PMC5967553.
- Mahdy OA, Nassar AM, Elsaywy BSM, Alzan HF, Kandil OM, Mahmoud MS, et al. Cross-sectional analysis of Piroplasma species-infecting camel (Camelus dromedaries) in Egypt using a multipronged molecular diagnostic approach. *Front Vet Sci.* 2023; 10: 1178511. doi: 10.3389/fvets.2023.1178511. PMID: 37187929; PMCID: PMC10175621.
- Oliveira-Sequeira TC, Oliveira MC, Araujo JP Jr, Amarante AF. PCR-based detection of Babesia bovis and Babesia bigemina in their natural host Boophilus microplus and cattle. *Int J Parasitol.* 2005; 35(1): 105-111. doi: 10.1016/j.ijpara.2004.09.002. PMID: 15619521.
- El-Sayed S, Rizk MA, Terkawi M, Igarashi I. Cocktail Babesia bovis antigens for global detection of Babesia bovis infection in cattle. *Exp Parasitol.* 2019; 206: 107758. doi: 10.1016/j.exppara.2019.107758. Epub 2019 Sep 12. PMID: 31521628.
- Eamens GJ, Bailey G, Jenkins C, Gonsalves JR. Significance of Theileria orientalis types in individual affected beef herds in New South Wales based on clinical, smear and PCR findings. *Vet Parasitol.* 2013; 196(1-2): 96-105. doi: 10.1016/j.vetpar.2012.12.059. Epub 2013 Jan 9. PMID: 23384579.
- Zhang XL, Deng YP, Yang T, Li LY, Cheng TY, Liu GH, Duan DY. Metagenomics of the midgut microbiome of Rhipicephalus microplus from China. *Parasit Vectors.* 2022; 15(1): 48. doi: 10.1186/s13071-022-05161-6. PMID: 35135613; PMCID: PMC8822867.
- Kumar R, Sharma AK, Ghosh S. Menace of acaricide resistance in cattle tick, Rhipicephalus microplus in India: Status and possible mitigation strategies. *Vet Parasitol.* 2020; 278: 108993. doi: 10.1016/j.vetpar.2019.108993. Epub 2019 Nov 26. PMID: 31954273.
- Makenov MT, Toure AH, Korneev MG, Sacko N, Porshakov AM, Yakovlev SA, et al. Rhipicephalus microplus and its vector-borne haemoparasites in Guinea: further species expansion in West Africa. *Parasitol Res.* 2021; 120(5): 1563-1570. doi: 10.1007/s00436-021-07122-x. Epub 2021 Mar 31. PMID: 33788020.
- Gómez GF, Isaza JP, Segura JA, Alzate JF, Gutiérrez LA. Metatranscriptomic virome assessment of Rhipicephalus microplus from Colombia. *Ticks Tick Borne Dis.* 2020; 11(5): 101426. doi: 10.1016/j.ttbdis.2020.101426. Epub 2020 Apr 4. PMID: 32473925.
- Ramos CM, Cooper SM, Holman PJ. Molecular and serologic evidence for Babesia bovis-like parasites in white-tailed deer (Odocoileus virginianus) in south Texas. *Vet Parasitol.* 2010; 172(3-4): 214-220. doi: 10.1016/j.vetpar.2010.05.004. Epub 2010 May 12. PMID: 20605333.