

Research Article

Effect of Elevated CO₂ and Low Temperature on Photosynthetic Potential of Indoor Plants

Ankur Agarwal*, Dinesh Pathak, Om Prakash, and Nasim M

Defence Institute of Bio Energy Research, India

*Corresponding author

Ankur Agarwal, Defence Institute of Bio Energy Research, Haldwani-263139, India, Tel: 091-5946-232800, Email: ankurdr@rediffmail.com

Submitted: 11 December 2017

Accepted: 28 December 2017

Published: 29 December 2017

ISSN: 2333-6668

Copyright

© 2017 Agarwal et al.

OPEN ACCESS

Keywords

- Temperature acclimation
- Cold tolerance
- Potted plants
- Air purifying plants
- Oxygen liberation during dark
- Transpiration rate
- Photosynthetic rate

Abstract

Several ornamental potted plant species have the ability to absorb pollutants and purify indoor air. The present study was aimed at to select best plant for pot transplantation in the cold areas of army installations. As diurnal variations in the temperature are high in these cold high altitude areas even in the indoor environment, plants were evaluated for their photosynthetic response under these conditions along with their oxygen releasing potential during light and dark conditions. Effect of temperature levels (10, 15, 20 and 25°C) and CO₂ levels (400, 420 and 450 ppm) were studied on net photosynthesis and transpiration rate in three indoor plants viz., spider plant (*Chlorophytum comosum*), dracaena (*Dracaena fragrans*) and snake plant (*Sansevieria trifasciata*). Decrease in photosynthetic as well as transpiration rate of these indoor plants was recorded with decrease in temperature from ambient (25°C) to 10°C. At 10°C, photosynthetic rate of Dracaena reduced to 1/7th of ambient whereas in case of snake and spider plants it was near to half only. Even though snake plant maintained the good photosynthetic rate (~5 $\mu\text{mol}/\text{m}^2/\text{s}$). Small increase in CO₂ levels was not detrimental to photosynthetic efficiency of indoor plants even at temperatures lower than ambient. At 10°C, snake plant exhibited the maximum transpiration rate (0.163 $\text{mmol}/\text{m}^2/\text{s}$). Among the three tested plants, Snake plant exhibited better oxygen releasing potential during both light and dark period.

INTRODUCTION

In the present century, current CO₂ levels are rising much rapidly than the expected rate which has attracted the attention of global community. Many studies have predicted an increase in the enhancement of carbon gain associated with elevated CO₂ at higher temperatures [1-3] and in general predict better performance of crop in an environment having elevated CO₂ combined with higher temperature due to global warming. Although crop specific response have also been reported showing better yield at lower temperatures under influence of higher carbon dioxide concentration [4].

Photosynthesis has long been recognized as one of the most temperature-sensitive processes in plants [3]. Photosynthesis is the primary process by which carbon enters the biosphere and by which plants sense rising CO₂ [5]. Photosynthesis is particularly sensitive to thermal stress, with increased photo inhibition of PSII observed at temperature extremes [6]. Increases in atmospheric levels of CO₂ above current levels can increase photosynthesis [1] even when plants were grown at temperature extremes [2]. Elevated CO₂ has been reported to increase water-use efficiency through regulation of stomatal conductance and transpiration [7]. Since plants face environmental conditions simultaneously,

interactive effect of temperature and CO₂ on photosynthesis is of paramount importance and has been review by Morison and Lawlor [2].

Indoor air quality has become a major issue in recent years. Volatile organic compounds (VOCs) in indoor air have received appreciable attention due to their adverse health effects on humans [8]. An alternative way to reduce the level of VOCs in indoor air is the use of plants. Several ornamental potted plant species have the ability to absorb VOCs [9] and purify indoor air [10,11]. Plants also offer the advantage of providing psychological and social benefits for humans [12]. The aim of the study was to select best plant for pot transplantation in the cold areas of army barracks to improve the indoor air quality because barracks are heated through keroheaters which is leading to poor indoor air quality. Since carbon dioxide levels are also generally high due to closed environment and poor air exchange rate during winter due to closed windows, effect of elevated carbon dioxide was also studied along with temperature. As diurnal variations in the temperature are high in these cold high altitude areas even in the indoor environment, plants were evaluated for their photosynthetic efficiency and oxygen releasing potential under light and dark conditions.

MATERIALS AND METHODS

Experimental materials and growing conditions

Fully grown potted-plants of three ornamental species were used in the present study. These were *Dracaena fragrans* (Dracaena), *Chlorophytum comosum* (Spider plant), *Sansevieria trifasciata* (Snake plant). All plants were exposed to four temperature regimes gradually from higher to lower (25°C, 20°C, 15°C and 10°C) and three CO₂ levels (400, 420 and 450 ppm) for 07 days in plant growth chamber LT-105 (Percival Scientific Inc., Perry, Iowa, USA) equipped with cold fluorescent light (25 K lux). Photoperiod of the growth chamber was adjusted to 12/12 h (day/night) and 70% humidity for all the treatments. Each treatment consisted of three pots of each species. Net photosynthesis was estimated by using a portable leaf chamber analyser (LCA-4, ADC Bio Scientific Ltd., UK). Data were collected from the all plants. Three samples were collected from each plant and average was worked out.

Estimation of oxygen releasing potential

To estimate the oxygen releasing potential, three selected potted plants viz., *Dracaena fragrans* (Dracaena), *Chlorophytum comosum* (Spider plant), *Sansevieria trifasciata* (Snake plant) were kept inside the leak proof air tight polythene bags of 150 gsm (Himedia) at ambient temperature. Mouth of the polythene was tied tightly with a provision of tubing to monitor the gas exchange with the help of multigas analyzer (GasAlertMicro5, BW Technologies by Honeywell, USA) as per schedule. Tubing was plugged suitably to avoid any leakage of gas. Plants were kept inside the polybags for 05 hrs with and without light to estimate the effect of light/dark conditions on oxygen releasing potential of these indoor plants. Dark conditions were created by covering the polythene bags with dark cotton cloth. Gas composition for oxygen content was checked on hourly basis. Plants selected were of uniform height and growth of one year age.

RESULTS

Effect on photosynthesis and transpiration rate

In the present experiment, results revealed that photosynthetic rates in all three plants decreased significantly as they were exposed to low temperature from ambient (Figure 1). This drop in photosynthetic rate was more pronounced for initial drop in temperature from 25 to 20°C and ranged from 50-58%. Whereas further decrease in temperature from 20 to 10°C resulted in to gradual decrease in photosynthetic rate except snake plant which exhibited higher photosynthetic rates compared to other two plants. Effect of higher CO₂ concentrations (>400 ppm) followed the same trend as of 400 ppm. The results also revealed that photosynthetic rate was marginally better at the highest CO₂ concentration (450 ppm) at ambient temperature but as the temperature dropped to 10°C from ambient, photosynthetic rates more or less dropped slightly at this CO₂ level. Corresponding decrease in photosynthetic rate of these indoor plants with decrease in temperature from ambient to 10°C exhibited the litmus paper test on the endurance of these plants. At 10°C, photosynthetic rate of Dracaena reduced to 1/7th of ambient whereas in case of snake and spider plants it was near to half only. Even though snake plant maintained the good

photosynthetic rate (~5 umol/m²/s).

Transpiration rate also decreased significantly with the decrease in temperature from ambient to 10°C (Figure 2) in all three plants. Average transpiration rate over all three CO₂ concentrations was comparatively higher in dracaena (0.564 mmol/m²/s) than rest two plants at ambient temperature whereas at 10°C, it was recorded maximum in snake plant (0.163 mmol/m²/s). Similarly to the photosynthetic rate, transpiration rate also exhibited steady decline during initial phase of temperature reduction and then decreased gradually except in snake plant which exhibited slightly increase in transpiration at 10°C at 400ppm CO₂ concentration. Variations in the transpiration rate due to change in CO₂ concentration were also non-significant at all temperature levels.




Figure 1 Effect of temperature (10, 15, 20 and 25°C) and CO₂ levels (400, 420 and 450 ppm) on net photosynthesis in three indoor plants; a) spider plant (*Chlorophytum comosum*), b) Dracaena (*Dracaena fragrans*) and c) Snake plant (*Sansevieria trifasciata*). Bar shown means (n = 3) ± SE are statistically significant (P<0.05) according to LSD test..

Effect on oxygen releasing potential

To estimate the oxygen releasing potential of these indoor plants, they were kept in air tight transparent polythene sheet. Gaseous oxygen composition of air inside the polythene bag was checked to measure the oxygen releasing potential of the indoor plants. Results revealed that all three plants behaved differently in their behavior of oxygen releasing potential [Figure 4, 5]. During the light sufficient conditions as presented in [Figure 4] decrease in gaseous oxygen content was rapid in dracaena during first hour whereas spider plant exhibited almost static oxygen content during first two hours and then gradual decrease in inside oxygen content in spider plant showing its better oxygen releasing potential. Snake plant exhibited a slight decrease in polythene gaseous content the first two hours but then it exhibited




Figure 2 Effect of temperature (10, 15, 20 and 25°C) and CO₂ levels (400, 420 and 450 ppm) on transpiration rate in three indoor plants; a) spider plant (*Chlorophytum comosum*), b) Dracaena (*Dracaena fragrans*) and c) Snake plant (*Sansevieria trifasciata*). Bar shown means ($n = 3$) \pm SE are statistically significant ($P < 0.05$) according to LSD test.




Figure 3 Concentration of Oxygen recorded inside the closed chamber to quantify the oxygen releasing potential of three indoor plants under a) light conditions, b) dark conditions. Bar shown means ($n = 3$) \pm SE are statistically significant ($P < 0.05$) according to LSD test.

positive trend in maintaining the oxygen content of gaseous content of polythene up to 5 hrs. During the dark conditions, oxygen releasing potential was also checked and results revealed that dracaena plant was poor in maintaining oxygen content inside the polythene during dark [Figure 5] whereas snake plant exhibited its potential in maintaining oxygen levels above 20.4% inside the polythene assembly even during dark conditions. Spider plants also exhibited its ability to cope with the dark and revealed its oxygen releasing potential even in dark conditions.

DISCUSSION AND CONCLUSIONS

Effect on photosynthesis and transpiration rate

The present study was basically aimed to study the response of indoor plants to low temperature ambience of cold region with elevated levels of CO₂ with study on interaction effect of elevated CO₂ and low temperature on plant response to photosynthetic and transpiration rate. The results in the present study revealed that photosynthetic rates in all three plants decreased significantly as they were exposed to low temperature from ambient. This drop in photosynthetic rate was more pronounced (50-58%) for initial drop in temperature from 25 to 20°C. Reduction in photosynthetic capacity, photochemical efficiency of PSII and apparent quantum yield has been reported under cold temperature conditions in potted plants compared to optimum temperature [13]. Slightly better net photosynthesis at 10°C temperature in snake plant compared to 15°C may be attributed to plant inherent capacity to adjust temperature change. Most plants show considerable

capacity to adjust their photosynthesis characteristics to their growing temperature by shifting temperature optimum required for photosynthesis and to counter the seasonal temperature shift [3] but ability of temperature acclimation of photosynthesis differ between plant species [14,15]. Snake plant has shown better cold tolerance potential in relation to photosynthetic ability [16]. The results also revealed that photosynthetic rate was marginally better at the highest CO₂ concentration (450 ppm) at ambient temperature but as the temperature dropped to 10°C from ambient, photosynthetic rates more or less dropped slightly at this CO₂ level. Although effect of CO₂ levels on net photosynthesis was somewhat similar at all temperatures, snake plant exhibited better performance. Higher photosynthesis under elevated CO₂ conditions in plants is largely due to lower rates of respiration and photorespiration [17]. When respiration is curtailed, carbon fixed during photosynthesis is available for anabolic process which may also result in increased photosynthetic capacity of plants [18, 19]. Increased levels of CO₂ result in closure of stomata in C₃ and C₄ plants which in turn increase water use efficiency and a consequence exhibit better photosynthetic efficiency [20, 21] as well as lower electron transport in C₃ and C₄ and CAM plants with decreased stomatal conductance [1]. At higher CO₂ concentration, water use efficiency increased in all temperatures due to increased canopy photosynthesis and decreased canopy respiration along with higher carbon assimilation [22] in cotton. Idso and Kimball [23] reported that doubling of CO₂ from 360 to 720 µl/l concentration doubled the net photosynthesis and decreased respiration by half in two tree species. A good correlation has been reported between the elevated CO₂ and increased level of chloroplast solutes [24,25] which may also have contributed to the phenomenon of higher photosynthesis.

Stomata opening are the common path for gas exchange and therefore transpiration is well correlated with photosynthesis. Transpiration regulates the water balance in plants and is influenced indirectly by environmental factors which affect stomatal opening [26]. In present study, transpiration rate also decreased significantly with the decrease in temperature from ambient to 10°C in all three plants. Average transpiration rate over all three CO₂ concentrations was comparatively higher in dracaena at ambient temperature whereas at low temperature (10°C), it was recorded maximum in snake plant, which in turn can be related to the better photosynthetic ability of snake plant at lower temperature. Transpiration was found to be temperature dependent and at sub optimal temperature for metabolic activity of *Vigna sinensis* transpiration was also low compared to optimal temperature [27]. Transpiration rates of plants have been found to decrease at higher CO₂ concentration despite increased leaf area index [28] which indicates greater canopy resistance to water vapour flux in high CO₂ environment but biomass increase in temperature dependent till achievement of maximum photosynthetic capacity [22].

Effect on oxygen releasing potential

Gaseous oxygen composition of air inside the polythene bag decreased with time in dracaena and spider plant but it remained somewhat static in snake plant in light as well dark conditions which indicated the good oxygen releasing potential of snake plant compared to other two. Srivatsan [29] also noted

that common house plants have a very unique pattern of O₂ production during the course of a day. All tested plants including snake plant attained peak of O₂ production during early morning hours much before sunlight reaches its maximum intensity. He also found that snake plant maintains good O₂ concentration even at the lower light intensities. Difference in diurnal gas exchange (O₂ and CO₂) has been observed between C₃ and C₄ plants species when photosynthetic photon flux density and water potential were moderate with better performance of C₄ [30]. Snake plant exhibited its potential in maintaining oxygen levels above 20.4% inside the polythene assembly in the present study. Snake plant, a CAM family plant, has exhibited better photosynthetic potential at low temperature compared to other indoor plants as well as tolerance to low temperature in our previous study [16]. Temperature dependent CO₂ fixation rate at day and night differ in CAM species plant Kalanchoe [3] although principal limiting step for CO₂ fixation at night was not stomatal conductance but malate formation by enzyme. Since photosynthesis and respiratory process are related, O₂ evolution depend on the balance of these two and positive relationship between these two indicates that higher amount of O₂ is released compared to its consumption at temperatures suitable for crop growth compared to lower or higher temperatures [31].

In conclusion, it can be said that small increase in CO₂ levels are not detrimental to photosynthetic efficiency of indoor plants even at temperatures lower than ambient but photosynthetic and transpiration rates of indoor plants decreases sharply at lower temperature. Although snake plant maintained almost double photosynthetic potential than dracaena including better oxygen releasing potential in light and dark conditions therefore could be a better candidate indoor plant for pot transplantation in cold locations to improve the indoor air quality as also have seen in our previous study [32].

REFERENCES

1. Wang D, Heckathorn SA, Barua D, Joshi P, Hamilton EW, Lacroix JJ. Effects of elevated CO₂ on the tolerance of photosynthesis to acute heat stress in C₃, C₄, and CAM species. *Am J Bot.* 2008; 95: 165-176.
2. Morison JIL, Lawlor DW. Interactions between increasing CO₂ concentration and temperature on plant growth, *Plant Cell Environ.* 1999; 22: 659-682.
3. Yamori W, Hikosaka K, Way DA. Temperature response of photosynthesis in C₃, C₄, and CAM plants: temperature acclimation and temperature adaptation. *Photosyn Res.* 2013; 2: 101-117.
4. Sun P, Mantri N, Lou H, Hu Y, Sun D, Tingting Dong, et al. Effects of elevated CO₂ and temperature on yield and fruit quality of strawberry at two levels of nitrogen application. *Plos One.* 2012; 7: e41000.
5. Drake BG, Gonzalez Meler MA, Long SP. More efficient plants: a consequence of rising atmospheric CO₂? *Ann Rev Plant Physiol Mol Biol.* 1997; 48: 609-639.
6. Weis E, Berry JA. Plants and high temperature stress. In: Long SP, Woodward JA editors. *Plants and temperature.* Cambridge: Company of Biologists. 1988; 329-346.
7. Ainsworth EA, Rogers A. The response of photosynthesis and stomatal conductance to rising [CO₂]: mechanisms and environmental interactions, *Plant Cell Environ.* 2007; 30: 258-270.
8. Jones AP. Indoor air quality and health. *Atmos Environ.* 1999; 33: 4535-4564.

9. Yang DS, Pennisi, Son KC, Kays SJ. Screening indoor plants for volatile organic pollutant removal efficiency, Hortscience. 2009; 44: 1377-1381.
10. Wolverton BC, Johanson A, Bounds K. Interior landscape plants for indoor air pollution abatement. NASA; 1989.
11. Cruz MD, Christensen JH, Thomsen JD, Muller. Can ornamental potted plants remove volatile organic compounds from indoor air? - A review. Environ Sci Pollut Res. 2014; 24: 13909-13928.
12. Bringslimark T, Hartig T, Patil GG. The psychological benefits of indoor plants: a critical review of the experimental literature. J Environ Psychol. 2009; 29: 422-433.
13. Oliveira G, Penuelas J. Effects of winter cold stress on photosynthesis and photochemical efficiency of PSII of the Mediterranean *Cistus albidus* L. and *Quercus ilex* L. Plant Eco. 2004; 175: 179-191.
14. Hill RS, Read J, Busby JR. The temperature-dependence of photosynthesis of some Australian temperate rainforest trees and its biogeographical significance. J Biogeogr. 1988; 15: 431-449.
15. Yamori W, Noguchi K, Hikosaka K, Terashima I. Phenotypic plasticity in photosynthetic temperature acclimation among crop species with different cold tolerances. Plant Physiol. 2010; 152: 388-399.
16. Gupta SM, Agarwal A, Dev B, Kumar K, Prakash O, Arya MC, Nasim M. Assessment of photosynthetic potential of indoor plants under cold stress. Photosynthetica. 2016; 54: 138-142.
17. Long SP, Drake BG. Photosynthetic CO₂ assimilation and rising atmospheric CO₂ Concentrations: In Crop Photosynthesis. Spatial and Temporal Determinants. N R Baker and H Thomas, Elsevier. Amsterdam: 1992; 69-95.
18. Devkumar AS, Shayee MSS, Udaykumar M, Prasad TG. Effect of elevated CO₂ concentration on seedling growth rate and photosynthesis in *Hevea brasiliensis*. J Biosci. 1988; 23: 33-36.
19. Makino A, Mae T, Photosynthesis and plant growth at elevated levels of CO₂, Plant Cell Physiol. 1999; 40: 999-1006.
20. Knapp AK, Hamerlyn CK, Owensby CE. Photosynthetic and water relations response to elevated CO₂ in the C₄ grass. *Andropogon gerardii*. Int J Plant Sc. 1993; 154: 459-466.
21. Ziska LH, Bunce JA. Influence of increasing carbon dioxide concentration on the photosynthetic and growth stimulation of selected C₄ crops and weeds. Photosyn Res. 1997; 54: 199-208.
22. Reddy VR, Reddy KR, Hodges HF. Carbon dioxide enrichment and temperature effects on cotton canopy photosynthesis, transpiration, and water-use efficiency. Field Crops Res. 1995; 41: 13-23.
23. Idso SB, Kimball BA. Effects of atmospheric CO₂ enhancement on net photosynthesis and dark respiration rates of three Australian tree species. J Plant Physiol. 1993; 141: 166-171.
24. Poorter H, Van Berkel Y, Baxter R, Den Hertog J, Dijkstra P, Gifeord RM et al. The effect of elevated CO₂ on the chemical composition and construction costs on leaves of 27 C₃ species. Plant Cell and Environment. 1997; 20: 472-482.
25. Taub DR, Seemann JR, Coleman JS. Growth in elevated CO₂ protects photosynthesis against high-temperature damage. Plant Cell and Environment. 2000; 23: 649-656.
26. Ku SB, Edwards GE, Tanner CB. Effects of light, carbon dioxide, and temperature on photosynthesis, oxygen inhibition of photosynthesis, and transpiration in *Solanum tuberosum*. Plant Physiol. 1977; 59: 868-872.
27. Neri D, Battistelli R, Albertini G. Effect of low light intensity and temperature on photosynthesis and transpiration of *Vigna sinensis* L. J. Fruit Ornamental Plant Res. 2003; 11: 17-24.
28. Reddy VR, Reddy KR, Acock B. Carbon dioxide and temperature effects on cotton leaf growth and development. Biotronics. 1994; 23: 59-74.
29. Srivatsan A. A Dual Sensor System for Determining the Unique Oxygen Production Signature of Plants. J. Selected Areas Mechatronics. 2005; 5: 1-5.
30. Niu SL, Jiang GM, Li YG, Gao LM, Liu MZ. Diurnal gas exchange and superior resources use efficiency of typical C₄ species in Hunshandak Sandland, China. Photosynthetica. 2003; 41: 221-226.
31. Ribeiro RV, Machado EC, de Oliveira RF. Temperature response of photosynthesis and its interaction with light intensity in sweet orange leaf discs under non-photorespiratory condition. Ciênc Agrotec, Lavras. 2006; 30: 670-678.
32. Agarwal A, Sabri M, Prakash O, Nasim M. Evaluation of in indoor plants for their pollution tolerance ability. JEAES. 2017; 5: 21-26.

Cite this article

Agarwal A, Pathak D, Prakash O, Nasim M (2017) Effect of Elevated CO₂ and Low Temperature on Photosynthetic Potential of Indoor Plants. Int J Plant Biol Res 5(4): 1078.