

## Review Article

# Review on Epidemiology and Public Health Importance of Zika Virus Disease

Gelma Debesa, Tedele Kebeta, and Wubishet Zewdie\*

Wollega University College of Medical and Health Sciences School of Veterinary Medicine  
Nekemte, Ethiopia

## \*Corresponding author

Wubishet Zewdie, Oromia Pastoral Area Development Commission Yabello Regional Veterinary Laboratory, P.O. Box 169, Yabello, Ethiopia, Email: wubevet1921@gmail.com; wubenbe2122@gmail.com

Submitted: 13 November 2018

Accepted: 26 November 2018

Published: 28 November 2018

## Copyright

© 2018 Zewdie et al.

## OPEN ACCESS

## Keywords

• Control; Transmission; Prevention; Zika virus; Zoonoses

## Abstract

We reviewed to document epidemiology and public health importance of Zika virus disease. The review was made by searching published paper online about Zika virus disease in different parts of the world. We noticed that the disease is zoonotic and economically important diseases as emerging diseases in different parts of the world. The disease seen and diagnosed in different parts of the world after first isolated from a rhesus monkey in the Zika forest of Uganda in 1947. In addition, serological evidence of infection was identified in a few animals in Indonesia, including horses, cows, goats and bats. The disease can appear anywhere; where these animals/reservoirs found and environment is favorable for mosquitoes breeding that are the primary vectors for virus in the world. Therefore, scientific communities, authorities and other concerned body need to give attention for the disease as it has public health and economic importance.

## INTRODUCTION

Zika virus (ZIKV) disease is caused by an arthropod-borne virus of family Flaviviridae genus *Flavivirus* [1]. It is an arbovirus which can cause diseases in humans through a mosquito vector that was daytime-active *A. aegypti* and *A. albopictus* mosquitoes. It was first isolated from rhesus monkey in the Zika forest in Uganda in 1947 [2]. The virus is most prevalent and emerging pathogen that has recently causing serious disease around the world. Cases of ZIKV disease were reported in Micronesia in 2007 and then in French Polynesia in 2013. In Brazil the virus was introduced and associated with cases of microcephaly [3]. The virus has the potential to spread anywhere due to day time active mosquitoes capability to spread this virus. There is no case report about Zika Virus disease in Ethiopia [4]. In non-human primates (NHP), antibodies against Zika Virus have been reported both in wild and experimental animals in Borneo [5] and Uganda [6].

There is no case report of Zika virus disease in Ethiopia, Even though some studies and literatures review have been conducted on the Zika virus disease, the viral pathogen is re-emerging and spread into different countries of the World. Therefore the objects of this review is to review the epidemiology and public health importance of Zika virus disease

## EPIDEMIOLOGY AND ZONOTIC IMPORTANCE OF ZIKA VIRUS DISEASE

### Etiology and morphology of the virus

Zika virus is a mosquito-borne disease caused by single strand, non-segmented, positive-sense RNA virus of family Flaviviridae genus *Flavivirus* (Figure 1). The virus infects and cause disease in humans and non-human primates (NHP) [7].

The structure of the Zika virus is similar to other Flaviviruses. The other complex structure of the genome is terminus which is essential for causing disease in human. The protein composes the majority of the virion surface and is involved in the viral-host cell binding and membrane fusion during replication [9].

### Geographical distributions and risk factor associated with diseases

The first virus isolation was case reported in human being in Uganda and Tanzania in 1954. Before 2007, there were reports of confirmed cases of Zika virus infection from the Africa and




Figure 1 Structure of Zika virus and its genome [8].

Southeast Asian continent. Later on in 2007, major epidemic occurred in Yap Island (Micronesia) where it affected about 5000 person [6], and also case report in Gabon [10].

In late 2015, the virus was reported for the first time in a number of countries in Central and South America with symptoms of microcephaly in affected parts of Brazil and now includes many Caribbean nations. However, the epicenter of the virus was Brazil and the virus soon spread to El Salvador, Honduras, Colombia, Venezuela and Costa Rica (Table 1).

The relationship of mosquito number with increased temperature is biological, that means it favors mosquitoes reproduction and increase their chance of contact with humans (for blood meal in case of female *A. aegypti*), which favor Zika transmission [22]. Risk factors associated with a disease are Vector related, Host related and Environmental related risk factors. The virus reservoir host is not completely identified and isolated, some studies suggests that, the reservoir host of Zika virus was primates. The other authors indicate the presence of anti-Zika antibodies in various animals including big mammals such as Zebras, Elephants and rodents in Pakistan [23]. The main source of Zika virus is infectious mosquitoes, mainly *Aedes* and secondary source is infected humans [24] (Figure 2).

Changes in climatic conditions, creates suitable conditions for mosquitoes by increasing its biting season and disease carrying mosquitoes will continue to bite as the climate warms. Weather conditions and global warming are important in occurrences and incidences of Zika Virus. For instance, El Niño is one of the factors that bring heavy rain which result in creating good opportunity for mosquito reproduction and multiplication site [25].

## Transmission

The virus transmissions can be based on the numbers of primates and mosquitoes vectors [26]. The most common route of Zika virus diseases transmission is through the bite of mosquitoes of the family Culicidae and genus *Aedes* (hematophagous arthropods) during their blood meal [24]. Other modes of transmission are non-vector borne transmission through sexual intercourse, blood transfusion, organ transplantation and prenatal transmission has been reported. Sexual transmission of virus is a




Figure 2 *A. albopictus* and *A. aegypti* [21].

particular concern during pregnancy. Transmission of the virus from infected women to their sex partners has not been reported. However, transmission from infected men to women partner has been reported [18]. Transmission is possible through organ transplantation, mother-to-fetus during the period of pregnancy [19]. People traveled from areas with active Zika virus disease transmission to countries with free of Zika virus can transmit the disease to their sex partners.

Transmission in laboratory form *A. aegypti* mosquitoes to mice and a monkey was reported [18,24].

## Pathogenesis

Zika virus has affinity towards human immature dendritic cells, dermal fibroblasts and epidermal keratinocytes. The virus may replicate in the midgut. Zika virus disease enters the cells through receptors that are found on the surface of skin and nerve cells. After entrance the virus replicates and distribute to muscles, heart, central nervous system (CNS), and to fetus by crossing the placental barrier through blood vessels and lymphatic [27]. Viral replication in the host cells causes the release of type I interferon [28]. Inside the cells, they use host mechanism and finally cause apoptosis and autophagy of the cells. It causes damage to the eye of infants [29].

Zika virus diseases use certain means to defeat the host defense of the trophoblast and by attacking neuronal tissues, it cause neuronal abnormalities in the fetus [30]. Some study show that, the virus have affinity towards brain cells which was demonstrated when intraperitoneal injection of virus into mice the virus cross the blood brain barrier. Neurons and glial cells were infected by virus and producing intracytoplasmic inclusions called viral factories. One recent study revealed that, Zika virus

Location	Countries reported	References
Africa	Senegal, Uganda, Nigeria, Ivory Coast, Gabon, Tanzania, Egypt Central African Republic Sierra Leone	[4,10-14]
Asia	Cambodia, India, Indonesia, Malaysia, Pakistan, Philippines, Singapore, Thailand, Japan and Vietnam,	[11,15-17]
Pacific	Micronesia/Yap, New Caledonia, south pacific, and Cook islands, French, Polynesia	[4,18,19]
Europe	Germany	[16]
America	Brazil, Barbados, Colombia, Ecuador, El Salvador, French Guiana, Guatemala, Guyana, Honduras, Martinique, Mexico, Panama, Paraguay, Puerto Saint Martin, Suriname, and Venezuela	[20] [21]

and man have a peptide in common which cause microcephaly and Guillain-Barré Syndrome (GBS) [31].

### Clinical signs

The virus has an incubation period of 3-12 days [32]. In confirmed case of Zika virus, clinical signs include acute fever, maculopapular skin rashes, non-purulent conjunctivitis, arthralgia, headache, myalgia and asthenia [33]. There are also less evident signs like anorexia, abdominal pain, vomiting, diarrhea, burning sensation of sole and palm [34; 35]. Symptoms are usually self-limiting and may last for 4-7 days [32].

### Diagnosis

Diagnostic testing for Flaviviruses is either direct detection of the virus RNA or detection of antibody responses to viral infection. In this manner virus can be detected from blood, other body fluids and tissues depending on stage of infection. For Zika virus, urine appears to have a higher viral detection than blood [36]. Reverse transcriptase-polymerase chain reaction (RT-PCR) methods used to detect both intact viral particles and replicating viral RNA inside cells. Serological tests for Zika virus on immunoglobulins (IgG and IgM) can be done in order to confirm the presence of neutralizing antibody infection [37].

### Treatment

Zika Virus disease is relatively mild and requires no specific treatment. People sick with this disease should get plenty of rest, drink enough fluids and treat pain and fever with common medicines like acetaminophen (paracetamol), and antihistamine for pruritic rash is necessary. Other non-steroidal anti-inflammatory drugs (NSAIDs) should be avoided until dengue can be ruled out to reduce the risk of hemorrhage [37]. It is also important to understand the link

### Prevention and control

Prevention and control relies on reducing mosquitoes through reduction of breeding sites and reducing contact between mosquitoes and people. Contact between mosquito and people can be prevented by using physical barriers such as window nets, closed doors and windows [6]. Integrated management of the *Aedes* mosquitoes through biological and chemical control is safe and cost effective. Integrated vector management systems are the most important prevention and control method [22]. In case of biological control, larvivorous fish are recommended for control of *Aedes aegypti* in large water bodies or large water Containers.

Endotoxin producing bacteria, *Bacillus thuringiensis* serotype H-14 has been an effective mosquito control agent [24].

Insecticide which can kill both adult and larvae are import control method in developing countries like Ethiopia. Larvicides include temephos and organophosphate compound. Adulticide recommend for the control of adult *Aedes* mosquito is Pyrethrum sprays [38].

### CONCLUSION

In conclusion, Zika virus disease is not a life threatening, but it is serious disease because of the birth defect. It is important to understand the link between microcephaly and the disease

because the link is used to develop vaccine. Therefore, People should aware about the transmission, prevention and control of diseases to avoid exposure of pregnant women to the vector and community water gathering around the house.

### REFERENCES

1. Faye O, Freire CC, Iamarino A, Faye O, de Oliveira JV, Diallo M, et al. Molecular evolution of Zika virus during its emergence in the 20(th) century. *PLoS Negl Trop Dis*. 2014; 8: e2636.
2. Document FAQ, Zika, Centre for emerging and zoonotic diseases, and division of public health surveillance and response. 2016; 12: 63-86.
3. Zanluca C, Melo VC, Mosimann AL, Santos GI, Santos CN, Luz K. First report of autochthonous transmission of Zika virus in Brazil. *Mem Inst Oswaldo Cruz*. 2015; 110: 569-572.
4. Hayes EB. Zika virus outside Africa. *Emerg Infect Dis*. 2009; 15: 1347-1350.
5. Wolfe ND, Kilbourn AM, Karesh WB, Rahman HA, Bosi EJ, Cropp BC, et al. Sylvatic transmission of arboviruses among Bornean orangutans. *Am J Trop Med Hyg*. 2001; 64: 310-316.
6. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med*. 2009; 360: 2536-2543.
7. Bowman LR, Runge-Ranzinger S, McCall PJ. Assessing the relationship between vector indices and dengue transmission: a systematic review of the evidence. *PLoS Negl Trop Dis*. 2014; 8: e2848.
8. Lindenbach BD, Rice CM. Molecular biology of flaviviruses. *Adv Virus Res*. 2003; 59: 23-61.
9. Clarke BD, Roby JA, Slonchak A, Khromykh AA. Functional non-coding RNAs derived from the flavivirus 3' untranslated region. *Virus Res*. 2015; 206: 53-61.
10. Grard G, Caron M, Mombo IM, Nkoghe D, Mboui Ondo S, Jiolle D, et al. Zika virus in Gabon (Central Africa)--2007: a new threat from *Aedes albopictus*? *PLoS Negl Trop Dis*. 2014; 8: e2681.
11. Robin Y, Mouchet J. Serological and entomological study on yellow fever in Sierra Leone. *Bull Soc Pathol Exot Filiales*. 1975; 68: 249-258.
12. Jan C, Languillat G, Renaudet J, Robin Y. [A serological survey of arboviruses in Gabon]. *Bull Soc Pathol Exot Filiales*. 1978; 7: 140-146.
13. Saluzzo JF, Gonzalez JP, Hervé JP, Georges AJ. Serological survey for the prevalence of certain arboviruses in the human population of the south-east area of Central African Republic (author's transl). *Bull Soc Pathol Exot Filiales*. 1981; 74: 490-499.
14. Diallo D, Sall AA, Diagne CT, Faye O, Faye O, Ba Y, et al. Zika virus emergence in mosquitoes in southeastern Senegal, 2011. *PLoS One*. 2014; 9: e109442.
15. Olson JG, Ksiazek TG, Suhandiman, Triwibowo. Zika virus, a cause of fever in Central Java, Indonesia. *Trans R Soc Trop Med Hyg*. 1981; 75: 389-393.
16. Tappe D, Rissland J, Gabriel M, Emmerich P, Gunther S. First case of laboratory-confirmed Zika virus infection imported into Europe, November 2013. *Euro Surveill*. 2014; 19: 20685.
17. Kutsuna S, Kato Y, Takasaki T, Moi M, Kotaki A, et al. Two cases of Zika fever imported from French Polynesia to Japan, December 2013 to January 2014 corrected. *Euro Surveill*. 2014; 19: 20683.
18. Musso D, Nhan T, Robin E, Roche C, Bierlaire D, Zisou K, et al. Potential for Zika virus transmission through blood transfusion demonstrated during an outbreak in French Polynesia, November 2013 to February 2014. *Euro Surveill*. 2014; 19: 20761.

19. Musso D, Roche C, Robin E, Nhan T, Teissier A, Cao-Lormeau VM. Potential sexual transmission of Zika virus. *Emerg Infect Dis*. 2015; 21: 359-361.
20. Marcondes CB, Ximenes MF. Zika virus in Brazil and the danger of infestation by *Aedes* (*Stegomyia*) mosquitoes. *Rev Soc Bras Med Trop*. 2015; 49: 1.
21. World Health Organization (WHO). Statement on the first meeting of the International Health Regulations (2005) Emergency Committee on Zikavirus and observed increase in neurological disorders and neonatal malformations. 2016.
22. Disease Control Manual - NL. (DCM) Vector-borne and Zoonotic Diseases. (DCM - NL, 2016). 2016.
23. Darwish MA, Hoogstraal H, Roberts TJ, Ahmed IP, Omar F. A sero-epidemiological survey for certain arboviruses (Togaviridae) in Pakistan. *Trans R Soc Trop Med Hyg*. 1983; 77: 442-445.
24. Petersen E, Wilson ME, Touch S, McCloskey B, Mwaba P, Bates M, et al. Rapid spread of Zika virus in the Americas - implications for public health preparedness for mass gatherings at the 2016 Brazil Olympic Games. *Int J Infect Dis*. 2016; 44: 11-15.
25. Alto BW, Juliano SA. Precipitation and temperature effects on populations of *Aedes albopictus* (Diptera: Culicidae), implications for range expansion. *J Med Entomol*. 2001; 38: 646-656.
26. Favoretto S, Araújo D, Oliveira D, Duarte N, Mesquita F, Zanotto P, et al. First detection of Zika virus in neotropical primates in Brazil, a possible new reservoir. 2016.
27. Chan JF, Choi GK, Yip CC, Cheng VC, Yuen KY. Zika fever and congenital Zika syndrome, an unexpected emerging arbovirus disease. *J Infect*. 2016; 72: 507-524.
28. Hamel R, Dejarnac O, Wichit S, Ekchariyawat P, Neyret A, Luplertlop N, et al. Biology of Zika virus infection in human skin cells. *J Virol*. 2015; 89: 8880-8896.
29. McCarthy M. Severe eye damage in infants with microcephaly is presumed to be due to Zika virus. *BMJ*. 2016; 352: 855.
30. Sadovsky Y, Clifton VL, Knöfler M. Editorial: Zika virus and Placenta. *Placenta*. 2016; 40.
31. Lucchese G, Kanduc D. Zika virus and autoimmunity, from microcephaly to Guillain-Barré syndrome, and beyond. *Autoimmun Rev*. 2016; 15: 801-808.
32. Barrera-Cruz A, Díaz-Ramos RD, López-Morales AB, Grajales-Muñiz C, Viniegra-Osorio A, Zaldívar-Cervera JA, et al. Technical guidelines for the prevention, diagnosis and treatment of Zika virus infection. *Rev Med Inst Mex Seguro Soc*. 2016; 54: 211-224.
33. Goeijenbier M, Slobbe L, van der Eijk A, de Mendonça Melo M, Koopmans MP, Reusken CB. Zika virus and the current outbreak, an overview. *Neth J Med*. 2016; 74: 104-109.
34. Basarab M, Bowman C, Aarons EJ, Cropley I. Zika virus. *BMJ*. 2016; 352: 1-7.
35. Korhonen EM, Huhtamo E, Smura T, Kallio-Kokko H, Raassina M, Vapalahti O. Zika virus infection in a traveler returning from the Maldives, June 2015. *Euro Surveill*. 2016; 21.
36. Bingham AM, Cone M, Mock V, Heberlein-Larson L, Stanek D, Blackmore C, et al. Comparison of test results for Zika virus RNA in urine, serum, and saliva specimens from persons with travel associated Zika virus disease—Florida, 2016. *MMWR Morb Mortal Wkly Rep*. 2016; 65: 475-478.
37. Centers for Disease Control and Prevention (CDC). Zika virus, clinical evaluation and disease. 2016.
38. Pan American Health Organization/World Health Organization. Epidemiological Update: Neurological syndrome, congenital anomalies and Zika virus infection. 2016.
39. Bicca-Marques JC. Outbreak of yellow fever affects howler monkeys in southern Brazil. *Lancet*. 2009; 15: 18-21.

## Cite this article

Debesa G, Kebeta T, Zewdie W (2018) Review on Epidemiology and Public Health Importance of Zika Virus Disease. *Ann Public Health Res* 5(3): 1081.