

Editorial

Sleep Disorders in Epileptic Patients: The Role of Epilepsy and the Role of AEDs

Verrotti A*, d'Alonzo R and Rinaldi VE

Department of Pediatrics, University of Perugia, Italy

*Corresponding author

Verrotti A, Department of Pediatrics, University of Perugia, Sant'andreadelleFratte; Fax: +39-075-5784415; Email: averrott@unich.it

Submitted: 07 July 2014

Accepted: 16 July 2014

Published: 18 July 2014

Copyright

© 2014 Verrotti et al.

OPEN ACCESS

Epilepsy and sleep disorders are major health problems and the prevalence of sleep disorders in children with epilepsy is under recognized. Childhood sleep disturbances originate from factors that are not closely related to the primary neurological disorder but are multifactorial in origin and include poor sleep hygiene, detrimental effects of seizures on sleep, effects of antiepileptic drugs and coexistence of primary sleep disorders or comorbid illness [1]. In general, children with epilepsy experience alterations in total sleep, sleep architecture, sleep latency and spontaneous arousals with a higher incidence of sleep fragmentation and daytime drowsiness [2]. In particular, NREM sleep is important for declarative memory consolidation whereas REM sleep is involved in non declarative and emotional memory [3]. Sleep related paroxysmal disorders in infancy and childhood represent a significant challenge for the clinicians with the distinction of nocturnal epilepsy from non epileptic sleep disorders [4]. The American Academy of Sleep Medicine (AASM) in 2005, elaborated an international classification (ICSD-2) dividing sleep disorders mimicking epilepsy in three groups: NREM and REM parasomnias, sleep related movement disorders and other paroxysmal nocturnal events [5]. Sleep has well-documented association with specific epilepsy syndromes. The most important amongst these are nocturnal frontal lobe epilepsy (FLE), benign epilepsy with centro-temporal spikes (BECTS), benign epilepsy of childhood with occipital paroxysms (BECOP), Landau-Kleffner syndrome (LKS), infantile spasms and electrical status epilepticus during slow wave sleep. The amount of baseline rhythmicity occurring in brain differs considerably between the states of sleep and wakefulness. It is therefore not surprising that various seizures types begin preferentially in specific sleep states [6]. The neurophysiologic process involved in the deepening of NREM sleep may also facilitate both seizures and interictal epileptiform discharges (IEDS). NREM sleep represents state of synchronization between the brainstem reticular activating system, thalamus and cortex. A progressive hyperpolarization of thalamo-cortical neurons is also determined by a reduction of the effect of acetylcholine which contributes to NREM sleep deepening [7]. In contrast, REM sleep tends to suppress IEDS and may restrict their field of distribution to the epileptogenic origin by inhibition of thalamo-cortical synchronizing mechanisms [8]. Seizures are most likely to occur during the descent to deeper levels of sleep. Patients with epilepsy generally have macrostructural

sleep abnormalities such as increased number and duration of awakenings during sleep, reduced sleep efficiency, reduced and fragmented REM sleep and increased stage shifts [7]. Among the causes of sleep fragmentation, sleep related breathing disorders can be considered a trigger for seizures and IEDS. This suggests that children with obstructive sleep apnea syndrome (OSAS) may have a dysfunction of the arousal system control which can be due to the effect of a primary brain insult as a predisposing factor for both OSAS and paroxysmal EEG activity [9]. Therefore treatment of these disorders may potentially improve seizure control. On the other hand antiepileptic drugs (AEDs) exert a beneficial action on seizures not only with direct effects on neuronal excitability but also by stabilizing sleep and reducing sleep transitions. Whether the improvement in sleep patterns is a direct consequence of the use of AEDs or the consequence of the suppression of epileptic manifestations is the object of many debates. The effects of AEDs on sleep have been studied independent of seizures showing both detrimental and beneficial effects. AEDs affect sleep architecture by inducing sedation or insomnia [10]. In addition some AEDs can be weight inducing, therefore fragmenting sleep [11]. Virtually all AEDs have effects on sleep architecture and many studies have analyzed these effects. In particular Class 1 studies in healthy adults suggest that phenobarbital and levetiracetam reduce REM sleep whereas it is enhanced by gabapentin [12-13]. Other studies demonstrated that clobazam decreases slow wave sleep in the same type of population while it is increased by levetiracetam, pregabalin and tiagabine. Clobazam also reduces sleep latency and arousals/wake time also reduced by levetiracetam, phenobarbital, tiagabine and pregabalin [14]. Class 3 evidence based studies show that carbamazepine favours sleep efficiency and extends total sleep time increasing slow wave sleep in healthy subjects and diminishes sleep latency, arousals REM sleep and wake time [15]. On the other hand class 1 evidence based studies in epileptic adult patients showed that pregabalin increases slow wave sleep that is instead reduced by levetiracetam [16]. Moreover, phenobarbital and gabapentin reduce sleep latency and arousals. Regarding slow wave sleep it has been seen that carbamazepine and gabapentin have an enhancing effect contrarily to ethosuximide. Conflicting results were seen with lamotrigine on REM sleep and slow wave sleep [17]. Phenobarbital and phenytoin cause a reduction in patients REM sleep that was not demonstrated with the administration

of ethosuximide or gabapentin. Finally class 3 evidence based studies show that phenobarbital increases daytime sleepiness, an effect that was not seen when topiramate, lamotrigine, zonisamide or vigabatrin were administered to epileptic patients [18]. There are very few studies concerning epileptic patients with problems initiating sleep. On the basis of previously performed studies a reduction of sleep latency in this population could be obtained by using phenytoin, phenobarbital, carbamazepine and clobazam. Another domain where further clinical trials are required is that of epileptic patients with difficulty maintaining sleep. Gabapentin, tiagabalin or pregabalin increasing N3 and having a positive effect on sleep consolidation could potentially achieve a good maintenance of sleep [19]. Another two aspects that could be explored are epileptic patients with insomnia or with excessive daytime sleepiness. In the first case these subjects could benefit from a therapy with carbamazepine, tiagabalin or pregabalin; on the other hand it is important to avoid the association of phenobarbital, valproic acid and levetiracetam in patients with daytime drowsiness.

The complicated and reciprocal relationship between sleep and epilepsy is a topic that has been intriguing physicians and researchers for many years. Whether sleep affects epilepsy or epilepsy modifies sleep has been extensively evaluated [20]. Very little literature exists on the mutual interaction of epilepsy, AEDs and sleep disorders in the pediatric population. Future research should explore possible therapeutic applications of antiepileptic drugs in epileptic patients presenting sleep comorbidities [7].

REFERENCES

1. Bazil CW. Epilepsy and sleep disturbance. *Epilepsy Behav.* 2003; 4 Suppl 2: S39-45.
2. Carney PR, Kohrman M H. Relation between epilepsy and sleep during infancy and childhood. In: Bazil CW Malow BA, Sammaritano MR, editors. *Sleep and epilepsy: the clinical spectrum.* Amsterdam: Elsevier science; 2002; 359-372.
3. Jain SV, Glauser TA. Effects of epilepsy treatments on sleep architecture and daytime sleepiness: an evidence based review of objective sleep metrics. *Epilepsia.* 2014; 55: 26-37.
4. Derry CP, Duncan JS, Berkovic SF. Paroxysmal motor disorders of sleep: the clinical spectrum and differentiation from epilepsy. *Epilepsia.* 2006; 47: 1775-1791.
5. American Academy of Sleep Medicine. International classification of sleep disorders, 2nd edition. In. *Diagnostic and coding manual.* Westchester (IL): American Academy of Sleep Medicine, 2005.
6. Bazil CW. Nocturnal seizures. *Semin Neurol.* 2004; 24: 293-300.
7. Kothare SV, Kaleyias J. Sleep and epilepsy in children and adolescents. *Sleep Med.* 2010; 11: 674-685.
8. Carotenuto M, Parisi P, Esposito M, Cortese S, Elia M. Sleep alterations in children with refractory epileptic encephalopathies: a polysomnographic study. *Epilepsy Behav.* 2014; 35: 50-53.
9. Miano S, Paolino MC, Peraita-Adrados R, Montesano M, Barberi S, Villa MP, et al. Prevalence of EEG paroxysmal activity in a population of children with obstructive sleep apnea syndrome. *Sleep.* 2009; 32: 522-529.
10. Romeo DM, Brogna C, Quintiliani M, Baranello G, Pagliano E, Casalino T, et al. Sleep disorders in children with cerebral palsy: neurodevelopmental and behavioral correlates. *Sleep Med.* 2014; 15: 213-218.
11. Kothare SV, Kaleyias J. The adverse effects of antiepileptic drugs in children. *Expert Opin Drug Saf.* 2007; 6: 251-265.
12. Karacan I, Orr W, Roth T, Kramer M, Thornby J, Bingham S, Kay D. Dose-related effects of phenobarbitone on human sleep-waking patterns. *Br J Clin Pharmacol.* 1981; 12: 303-313.
13. Cicolin A, Magliola U, Giordano A, Terreni A, Bucca C, Mutani R. Effects of levetiracetam on nocturnal sleep and daytime vigilance in healthy volunteers. *Epilepsia.* 2006; 47: 82-85.
14. Nicholson AN, Stone BM, Clarke CH. Effect of the 1,5-benzodiazepines, clobazam and triflurazepam, on sleep in man. *Br J Clin Pharmacol.* 1977; 4: 567-572.
15. Riemann D, Gann H, Hohagen F, Bahro M, Müller WE, Berger M. The effect of carbamazepine on endocrine and sleep EEG variables in a patient with 48-hour rapid cycling, and healthy controls. *Neuropsychobiology.* 1993; 27: 163-170.
16. de Haas S, Otte A, de Weerd A, van Erp G, Cohen A, van Gerven J, et al. Exploratory polysomnographic evaluation of pregabalin on sleep disturbance in patients with epilepsy. *J Clin Sleep Med.* 2007; 3: 473-478.
17. Bonanni E, Galli R, Maestri M, Pizzanelli C, Fabbrini M, Manca ML, Ludice A. Daytime sleepiness in epilepsy patients receiving topiramate monotherapy. *Epilepsia.* 2004; 45: 333-337.
18. Wolf P, Röder-Wanner UU, Brede M. Influence of therapeutic phenobarbital and phenytoin medication on the polygraphic sleep of patients with epilepsy. *Epilepsia.* 1984; 25: 467-475.
19. Placidi F, Diomedes M, Scalise A, Marciani MG, Romigi A, Gigli GL, et al. Effect of anticonvulsants on nocturnal sleep in epilepsy. *Neurology.* 2000; 54: S25-32.
20. Méndez M, Radtke RA. Interactions between sleep and epilepsy. *J Clin Neurophysiol.* 2001; 18: 106-127.

Cite this article

Verrotti A, d'Alonzo R, Rinaldi VE (2014) Sleep Disorders in Epileptic Patients: The Role of Epilepsy and the Role of AEDs. *J Sleep Med Disord* 1(1): 1001.