

Research Article

The Outcomes of Female Soccer Players Undergoing ACL Reconstruction with Quadriceps Tendon Vs Patellar Tendon Autografts

Zachary Herman* and Dharmesh Vyas

Department of Orthopedic Surgery, PGY3, Orthopedic Surgery Resident, USA

***Corresponding author**

Zachary Herman, Department of Orthopedic Surgery, PGY3, Orthopedic Surgery Resident, USA, Tel: 7245162858

Submitted: 08 July 2023

Accepted: 30 August 2023

Published: 31 August 2023

ISSN: 2379-0571

Copyright

© 2023 Herman Z, et al.

OPEN ACCESS

Abstract

Purpose: The purpose is to compare functional outcomes, return to soccer rates, and revision rates in an all-female soccer player cohort undergoing quadriceps tendon (QT) autograft ACLR versus bone-patellar tendon-bone (BPTB) autograft ACLR.

Methods: Female soccer players who sustained an ACL rupture and underwent primary anatomic, single-bundle ACLR with BPTB autograft or QT autograft were included. Demographic and surgical characteristics were collected. Outcomes of interest included Tegner score, International Knee Documentation Committee (IKDC) score, Marx score, return to soccer rates, and failure rates.

Results: Data on 23 patients undergoing BPTB autograft ACLR and 14 undergoing QT autograft ACLR was available. Average age was 18.7 years, and average follow up was 4.8 years. Overall, 76% (28/37) returned to soccer and 5.4% (2/37) underwent revision ACLR. No major significant differences were found in demographic or surgical characteristics. No differences were found in postoperative IKDC scores, preoperative, postoperative, or change from pre- to postoperative Marx activity scores, or pre- and postoperative Tegner scores between the groups. QT autograft ACLR patients had significantly less change in Tegner scores pre- to postoperatively compared to the BPTB autograft ACLR group (0.6 + 1.2 versus 2.1 + 1.8; $p = 0.02$). Both groups had similar rates of return to soccer [78% (18/23) BPTB autograft ACLR versus 71% (10/14) QT autograft ACLR; $p = 0.64$].

Conclusion: BPTB autograft ACLR and QT autograft ACLR produced comparable, successful functional and return to soccer outcomes in this all-female soccer player cohort study. Larger, prospective studies are needed to improve the strength of conclusions and provide more information on the optimal graft choice for female soccer players.

Study Design: Cohort Study, Level III

Clinical Relevance: Surgeons can use the results of this study to counsel female soccer players on graft options and expected outcomes after ACLR with QT or BPTB autografts.

INTRODUCTION

Female athletes have been reported as two to eight times more likely to experience ACL injuries than male athletes [1,2], and in particular, female soccer players are nearly three times more likely than male soccer players to injure their ACL [3]. Furthermore, female soccer players are at a high risk of failure after ACL reconstruction (ACLR) upon returning to the field [4]. A 2021 study found these athletes had a two to five times higher risk of ipsilateral knee injury and knee surgery compared to healthy controls and female athletes who did not return to play following ACLR [5].

Due to the high risk of failure after ACLR in the female soccer

player population, there remains a debate on the optimal graft choice in these patients. A 2020 retrospective study compared outcomes of hamstring autograft and bone-patellar tendon-bone (BPTB) autograft ACLR in female soccer players and found comparable patient reported outcomes between the groups with similar rates of return to soccer [6]. Yet, hamstring autograft may be decreasing in popularity in the female athlete, as other studies have shown higher risk of failure in this population [7,8]. Furthermore, given results supporting the quadriceps tendon (QT) autograft's decreased donor site morbidity [9] as well as its proven success in functional knee recovery and limiting failure, it is becoming a more common choice in adolescent athletes [10-12]. As such, its results should be further examined in the female soccer player population.

Currently, limited data exists comparing outcomes of female soccer players after ACLR with QT autograft to other graft options. As such, the purpose of this study was to compare functional and return to soccer outcomes as well as operative knee reinjury and revision ACLR rates between BPTB autograft and QT autograft ACLR in an all-female soccer player cohort. We hypothesized that both graft types would result in similar functional outcomes, return to soccer rates, and revision rates in female soccer players.

METHODS

Data Collection

This combined retrospective study was approved by the Institutional Review Board at the University of Pittsburgh, IRB#: STUDY22050138 and IRB#: STUDY19030196. All female soccer players who sustained an ACL rupture underwent primary anatomic, single-bundle ACLR with BPTB autograft or QT autograft between 2015 and 2021 by one of seven sports medicine fellowship trained orthopaedic surgeons were included for analysis. Exclusion criteria included males, patients <14 years old, revision ACLR, double-bundle ACLR, multi-ligament knee reconstruction, all-epiphyseal techniques, and follow-up length less than 2 years. Demographic data including patient age, body mass index (BMI), level of soccer competition [defined as high school/recreational or collegiate (no professionals included in study)], laterality of injury, time to surgery from injury, and completion of physical rehabilitation with a licensed physical therapist prior to surgery were collected. Surgical characteristics including regional nerve block administration, procedure time, and concomitant meniscal procedure were recorded. All patients included for analysis were prospectively contacted for completion of a questionnaire assessing postoperative International Knee Documentation Committee (IKDC) [13,14] score, pre- and postoperative Marx activity scale scores [15], and pre-and postoperative Tegner scores [16], return to soccer information (including reason for failure to return), future ipsilateral knee injury (subjectively defined by patients completing the questionnaire), and ipsilateral revision ACLR. The primary outcome of interest was Tegner score. Secondary outcomes of interest included IKDC, Marx score, RTS rates, and failure rates. All questionnaires were completed using an institutional REDCap database (REDCap, Vanderbilt University) and stored alongside retrospective data collection. Follow-up length was determined using date of questionnaire completion.

Power Analysis

An a priori power analysis was conducted utilizing data from a previous manuscript investigating differences in outcomes between female soccer players undergoing BPTB and hamstring ACLR [6]. It was found that 17 and 25 individuals were required for the quad tendon and BPTB groups, respectively, to observe a difference of 1.8 points in Tegner scores with a power of 0.8 (alpha=0.05, beta=0.2).

Statistical Analysis

All data was analyzed using Microsoft Excel (V16.6) and SPSS V28 (IBM). Descriptive statistics were reported for all

demographic variables, surgical characteristics, and post-operative outcomes and displayed as means with standard deviations (SD) or numbers (n) with proportions (%). Continuous data was compared between BPTB autograft ACLR and QT autograft ACLR groups with independent t-test or Mann-Whitney U test for parametric and non-parametric data, respectively. Categorical data was compared between BPTB autograft ACLR and QT autograft ACLR groups with chi-squared tests. Statistical significance was set at p<0.05.

RESULTS

A total of 123 consecutive female soccer players met inclusion for the study, 53 undergoing QT autograft ACLR and 70 undergoing BPTB ACLR. Nine (9) patients were excluded from the BPTB autograft ACLR group as they underwent revision ACLR with BPTB autograft (hamstring autograft for primary ACLR). After survey distribution under protocol derived by the Institutional Review Board and completion by participants, data on 37 total female soccer players were available: 23 patients undergoing BPTB autograft ACLR and 14 undergoing QT autograft ACLR. Table 1 shows data on the entire study population. Average age was 18.7 years, and average follow up was 4.8 years. Overall, 76% (28/37) returned to soccer and 5.4% (2/37) underwent revision ACLR.

Table 2 displays a comparison of demographic and surgical

Table 1: Demographic characteristics of study population

Variable	Total Cohort (n = 37)
Age (years), mean (SD)	18.7 (4.0)
BMI, mean (SD)	22.0 (2.7)
Laterality (right), n (%)	15 (40.5)
Follow-Up Length (years), mean (SD)	4.8 (1.3)
Return to soccer (yes), n (%)	28 (75.7)
Return to soccer at higher level*, n (%)	9 (32.1)
Return to soccer at same level*, n (%)	13 (46.4)
Return to soccer at lower level*, n (%)	6 (21.4)
Revision ACLR	2 (5.4)

SD = standard deviation, n = number of patients, BMI = body mass index, * Indicates percentage of patients among those who returned to soccer

Table 2: Comparison of demographic and surgical characteristics between BPTB autograft ACLR versus QT autograft ACLR

Variable	BPTB (n = 23)	QT (n = 14)	p-value ^a
Age (years), mean (SD)	18.1 (2.6)	19.7 (5.4)	0.96
BMI, mean (SD)	21.7 (2.1)	22.3 (3.4)	1.00
Laterality (right), n (%)	9 (39.1)	6 (42.9)	0.82
Level of Competition* (College), n (%)	10 (43.5)	2 (14.3)	0.07
Completed Prehab (yes), n (%)	7 (30.4)	7 (50.0)	0.23
Time to Surgery (months), mean (SD)	1.1 (0.7)	3.7 (6.0)	0.03
Regional Nerve Block (yes), n (%)	17 (73.9)	9 (64.3)	0.40
Concomitant Meniscal Procedure (yes), n (%)	12 (52.2)	8 (57.1)	0.77
Procedure Time (minutes), mean (SD)	111.5 (29.1)	119.0 (26.9)	0.70
Follow-Up Length (years), mean (SD)	5.1 (1.4)	4.5 (1.0)	0.36

BMI = body mass index, SD = standard deviation, n = number of patients, BPTB = bone-patellar tendon-bone, QT = quadriceps tendon, ACLR = ACL reconstruction
* Level of competition: College versus high school/recreational
^aStatistical significance at p < 0.05 (bold).

characteristics between patients undergoing BPTB autograft ACLR versus QT autograft ACLR. No significant differences were found between age at surgery, BMI, laterality of injured knee, level of competition, or completion of rehabilitation prior to surgery. Mean time from injury to surgery was longer in the QT autograft ACLR group (3.7 ± 6.0 months versus 1.0 ± 0.7 ; $p = 0.03$). One patient in the QT autograft ACLR group had a time from injury to surgery of over 24 months, likely contributing to the wide standard deviation in the QT autograft ACLR group and this statistical finding. In terms of surgical characteristics, no significant differences were found between the groups with respect to regional nerve block usage, procedure time, and or percentage of female athletes undergoing concomitant meniscal procedure. Lastly, mean follow up was comparable between the groups, as the BPTB autograft group had average follow up of 5.1 years, and the QT autograft group had average follow up of 4.5 years ($p = 0.36$).

A comparison of outcomes of BPTB autograft ACLR versus QT autograft ACLR is shown in Table 3. No differences were found in postoperative IKDC scores, preoperative, postoperative, or change from pre- to postoperative Marx activity scores, or pre-and postoperative Tegner scores between the groups. QT autograft ACLR patients had significantly less change in Tegner scores pre- to postoperatively compared to the BPTB autograft ACLR group (0.6 ± 1.2 versus 2.1 ± 1.8 ; $p = 0.02$). Both groups had similar rates of return to soccer [78% (18/23) BPTB autograft ACLR versus 71% (10/14) QT autograft ACLR], subsequent ipsilateral knee injury [30% (7/23) BPTB autograft ACLR versus 21% (3/14) QT autograft ACLR], and revision ACLR [9% (2/23) BPTB autograft ACLR versus 0% (0/14) QT autograft ACLR].

Lastly, female soccer players who did not return to soccer after ACLR were asked to choose as many answers that applied from a list of reasons as to why they did not return. These results are shown in Table 4. Twenty-one percent (22%; 5/23) of patients undergoing BPTB autograft ACLR while 29% (4/14) patients undergoing QT autograft ACLR did not return to soccer.

Table 3: Comparison of patient-reported outcomes between BPTB autograft ACLR versus QT autograft ACLR

Variable	BPTB (n = 23)	QT (n = 14)	p-value ^a
Post-Operative IKDC, mean (SD)	85.8 (11.20)	91.6 (9.7)	0.10
Pre-Operative Marx, mean (SD)	14.7 (3.1)	14.6 (1.8)	0.51
Post-Operative Marx, mean (SD)	11.0 (5.3)	11.8 (4.6)	0.75
Change in Pre- to Post-Op Marx, mean (SD)	3.7 (4.3)	2.9 (3.8)	0.70
Pre-Operative Tegner, mean (SD)	9.4 (1.1)	8.9 (1.0)	0.12
Post-Operative Tegner, mean (SD)	7.3 (2.3)	8.2 (1.6)	0.23
Change in Pre- to Post-Op Tegner, mean (SD)	2.1 (1.8)	0.6 (1.2)	0.02
Return to soccer (yes), n (%)	18 (78.3)	10 (71.4)	0.64
Subsequent Ipsilateral Knee Injury (yes), n (%)	7 (30.4)	3 (21.4)	0.64
Subsequent Revision ACLR (yes), n (%)	2 (8.7)	0 (0)	0.26

IKDC = International Knee Documentation Committee, BPTB = bone-patellar tendon-bone, QT = quadriceps tendon, ACLR = ACL reconstruction, SD = standard deviation, n = number of patients
^aStatistical significance at $p < 0.05$ (bold).

Table 4: Descriptive comparison of reasons for failing to return to soccer BPTB autograft ACLR versus QT autograft ACLR

Reason for Failure to Return to Soccer	BPTB (n = 5)	QT (n = 4)
Not Cleared to Play, n (%)	0 (0)	0 (0)
Fearful of Re-Injury or Lack of Confidence in Knee, n (%)	3 (60)	2 (50)
Interests Have Changed, n (%)	1 (20)	2 (50)
Ipsilateral Re-Injury or Complication, n (%)	2 (40)	0 (0)
Contralateral Injury or Complication, n (%)	0 (0)	0 (0)
Season not Started or Sport not Available, n (%)	1 (20)	0 (0)
Other, n (%)	2 (40)	2 (50)

n = number of patients, BPTB = bone-patellar tendon-bone, QT = quadriceps tendon

Out of the reasons listed, 60% (3/5) of those undergoing BPTB autograft ACLR and 50% (2/4) of those undergoing QT autograft ACLR selected that “fear of re-injury or lack of confidence in the knee” kept them from returning. Forty percent (40%; 2/5) patients undergoing BPTB autograft ACLR selected “ipsilateral re-injury or complication” as a reason for preventing them from returning to soccer. Other reasons for not returning to soccer for athletes in both groups included “interests have changed”, “the season has not started yet or the sport is no longer available”, and “other”. Full results are listed in Table 4.

DISCUSSION

The main findings of this study were that both BPTB autograft ACLR and QT autograft ACLR result in similar, successful patient reported outcomes in a unique study population of all female soccer players with high return to soccer rates of 78.1% after BPTB autograft ACLR and 71.4% after QT autograft ACLR and low failure rates of 9% after BPTB autograft ACLR and 0% after QT autograft ACLR.

Functional knee outcomes in this study are comparable with the literature. A 2020 retrospective study with average follow up of 3.4 years comparing all-female soccer player cohorts undergoing BPTB autograft ACLR versus hamstring autograft ACLR found mean postoperative Tegner scores of 6.0 in the 41 athletes undergoing BPTB autograft ACLR [6]. While these results were statistically superior to the hamstring autograft ACLR cohort (mean postoperative Tegner 4.2) [6], postoperative Tegner scores for the BPTB autograft ACLR group in the 2020 study are comparable to the mean postoperative Tegner scores for both cohorts in this study (7.3 BPTB autograft ACLR; 8.2 QT autograft ACLR). Interestingly, in this study, mean change from pre- to postoperative Tegner scores were significantly lower in the QT autograft group (0.6) compared to the BPTB autograft group (2.1), suggesting that patients undergoing QT autograft ACLR may have retained a higher level of activity at follow up than those undergoing BPTB autograft ACLR. Furthermore, postoperative IKDC scores were similar between cohorts, with mean scores of 85.8 and 91.6 for the BPTB autograft ACLR and QT autograft ACLR groups respectively, and both cohorts' mean IKDC values reached the patient acceptable symptom state (PASS) of 75.9 defined by a 2016 study [17]. Overall, the results of patient reported outcomes in this study support the literature on a wider scale of patient populations, as data has shown the

effectiveness of both BPTB and QT autografts in functional knee recovery after ACLR in athletes [18-20].

Return to soccer rate in the entire study population was 76% (28/37), and when broken down by graft type, both cohorts had similar rates of return [78% (18/23) BPTB autograft ACLR versus 71% (10/14)] QT autograft ACLR. These results are higher than rates reported in the literature, as two case series of female soccer players from 2015 and a 2020 study comparing BPTB autograft ACLR versus hamstring autograft ACLR in female soccer players showed return to soccer rates between 46%-66% at similar follow up timepoints [6,21,22]. Recent data has focused on reasoning for failing to return to sport after ACLR. Fear or lack of psychological readiness is commonly cited as a reason for failed or delayed return to sport [6,23,24], and some studies show that females may be slower to develop psychological readiness to return to sport than males [25,26]. In this study, despite small sample sizes and small percentage of patients failing to return to soccer, "fear of reinjury and lack of confidence in operative knee" was still in the majority of reasons chosen for lack of return. As such, a focus of the postoperative rehabilitation should continue to include interventions designed to increase confidence and psychological readiness such as visualization of success, routine assessment of self-efficacy and fear of reinjury, strict adherence to physical therapy programs, goal-setting, and positive self-talk [27-29].

Female soccer players are at higher risk of failure after ACLR compared to non-soccer female athletes undergoing ACLR [4]. A 2016 cohort study reviewed medical records of female athletes undergoing ACLR, dividing patients into 2 groups: female soccer athletes and matched female athletes that did not participate in soccer [4]. Over an average follow up time of 68.8 months, the authors found that soccer players had higher rates of ACLR failure compared to non-soccer athletes (11% vs 1%) [4]. Furthermore, in a comparative study of BPTB autograft ACLR versus hamstring autograft ACLR in an all-female soccer player population, overall failure rate was reported as 11.3% (8/71), with 9.8% (4/41) failure rate for the BPTB autograft ACLR group and 13.3% (4/30) for the hamstring autograft ACLR group [6]. While approximately 27% (10/37) of female soccer players in this study reported a subsequent injury to ipsilateral knee, a smaller percentage experienced a retear requiring revision ACLR [5.4% (2/37) overall; 8.7% (2/23) BPTB autograft ACLR; 0% (0/14) QT autograft ACLR]. Despite the long duration of follow up, these low revision rates may be due to the small sample size of the study, relying on a survey to assess the status after ACLR and placing the study at risk for sampling bias. It is also possible that, as mean Tegner and Marx scores have decreased pre- to postoperatively, some of the athletes no longer participate in sport, resulting in decreased strenuous activity and less stress on their ACL graft.

The literature has shown that female soccer players that have undergone ACLR have almost a five-fold higher incidence of developing new knee-related injuries (ipsilateral ACL retear, contralateral ACL injury, other non-ACL related injury) than healthy-knee controls [30]. It is worth questioning why these

athletes continue to fail and reinjure, especially at higher rates than other athletic populations [30]. As the optimal graft type for the female soccer player continues to be studied, focus on other factors such as multicomponent exercise-based prevention programs, coronal and sagittal malalignment, rotatory instability, and postoperative neuromuscular training as well as quadriceps, hamstring, and core muscle development should continue in this population in attempt to provide these athletes with successful return to the field and the lowest possible chance of reinjury [31-33].

Despite its uniqueness in presenting outcomes of QT autograft ACLR in an all-female soccer player population, several limitations to this study exist and include its retrospective nature and low sample size, which is underpowered according to an a priori analysis performed. This study also relied on survey responses to address current knee function as well as rates of return to soccer and ACLR revision, introducing the possibility for sampling bias. Larger, randomized studies are needed to draw firmer conclusions on graft types with the most successful outcomes following ACLR in female soccer players.

CONCLUSIONS

BPTB autograft ACLR and QT autograft ACLR produced comparable, successful functional and return to soccer outcomes in this all-female soccer player cohort study. Larger, prospective studies are needed to improve the strength of conclusions and provide more information on the optimal graft choice for female soccer players. Surgeons can use the results of this study to counsel female soccer players on expected outcomes after ACLR.

REFERENCES

- Toth AP, Cordasco FA. Anterior cruciate ligament injuries in the female athlete. *J Gend Specif Med.* 2001;4:25-34.
- Arendt E, Dick R. Knee injury patterns among men and women in collegiate basketball and soccer. NCAA data and review of literature. *Am J Sports Med.* 1995; 23: 694-701.
- Prodromos CC, Han Y, Rogowski J, Joyce B, Shi K. A meta-analysis of the incidence of anterior cruciate ligament tears as a function of gender, sport, and a knee injury-reduction regimen. *Arthroscopy.* 2007;23:1320-1325.
- Allen MM, Pareek A, Krych AJ, Hewett TE, Levy BA, Stuart MJ, et al. Are Female Soccer Players at an Increased Risk of Second Anterior Cruciate Ligament Injury Compared With Their Athletic Peers? *Am J Sports Med.* 2016; 44: 2492-2498.
- Fältström A, Kvist J, Häggglund M. High Risk of New Knee Injuries in Female Soccer Players After Primary Anterior Cruciate Ligament Reconstruction at 5- to 10-Year Follow-up. *Am J Sports Med.* 2021; 49: 3479-3487.
- Britt E, Ouillette R, Edmonds E, Chambers H, Johnson K, Bastrom T, et al. The Challenges of Treating Female Soccer Players With ACL Injuries: Hamstring Versus Bone-Patellar Tendon-Bone Autograft. *Orthop J Sports Med.* 2020; 8: 2325967120964884.
- Tiplady A, Love H, Young SW, Frampton CM. Comparative Study of ACL Reconstruction With Hamstring Versus Patellar Tendon Graft in Young Women: A Cohort Study From the New Zealand ACL Registry. *Am J Sports Med.* 2023; 51: 627-633.

8. Salem HS, Varzhapetyan V, Patel N, Dodson CC, Tjoumakaris FP, Freedman KB. Anterior Cruciate Ligament Reconstruction in Young Female Athletes: Patellar Versus Hamstring Tendon Autografts. *Am J Sports Med.* 2019;47:2086-2092.
9. Mouarbes D, Dagneau L, Olivier M, Lavoue V, Peque E, Berard E, et al. Lower donor-site morbidity using QT autografts for ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2020; 28: 2558-2566.
10. Arnold MP, Calcei JG, Vogel N, Magnussen RA, Clatworthy M, Spalding T, et al. ACL Study Group survey reveals the evolution of anterior cruciate ligament reconstruction graft choice over the past three decades. *Knee Surg Sports Traumatol Arthrosc.* 2021; 29: 3871-3876.
11. Vaughn NH, Dunleavy ML, Jackson T, Hennrikus W. The outcomes of quadriceps tendon autograft for anterior cruciate ligament reconstruction in adolescent athletes: a retrospective case series. *Eur J Orthop Surg Traumatol.* 2022; 32: 739-744.
12. Brinkman J, Makovicka J, Economopoulos K. Paper 02: ACL Reconstruction in Cutting and Pivoting Athletes: Soft Tissue Quadriceps Tendon Autograft vs. Bone-Patellar Tendon-Bone Autograft. *Orthop J Sports Med.* 2022; 10: 2325967121S00566.
13. Irrgang JJ, Anderson AF, Boland AL, Harner CD, Kurosaka M, Neyret P, et al. Development and validation of the international knee documentation committee subjective knee form. *Am J Sports Med.* 2001; 29: 600-613.
14. Irrgang JJ, Ho H, Harner CD, Fu FH. Use of the International Knee Documentation Committee guidelines to assess outcome following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 1998; 6: 107-114.
15. Marx RG, Stump TJ, Jones EC, Wickiewicz TL, Warren RF. Development and evaluation of an activity rating scale for disorders of the knee. *Am J Sports Med.* 2001; 29: 213-218.
16. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res.* 1985: 43-49.
17. Muller B, Yabroudi MA, Lynch A, Chung-Liang Lai, van Dijk CN, H Fu F, et al. Defining Thresholds for the Patient Acceptable Symptom State for the IKDC Subjective Knee Form and KOOS for Patients Who Underwent ACL Reconstruction. *Am J Sports Med.* 2016; 44: 2820-2826.
18. DeFazio MW, Curry EJ, Gustin MJ, Sing DC, Abdul-Rassoul H, Ma R, et al. Return to Sport After ACL Reconstruction With a BTB Versus Hamstring Tendon Autograft: A Systematic Review and Meta-analysis. *Orthop J Sports Med.* 2020; 8: 2325967120964919.
19. Mouarbes D, Menetrey J, Marot V, Courtot L, Berard E, Cavaignac E. Anterior Cruciate Ligament Reconstruction: A Systematic Review and Meta-analysis of Outcomes for Quadriceps Tendon Autograft Versus Bone-Patellar Tendon-Bone and Hamstring-Tendon Autografts. *Am J Sports Med.* 2019; 47: 3531-3540.
20. Dai W, Leng X, Wang J, Cheng J, Hu X, Ao Y. Quadriceps Tendon Autograft Versus Bone-Patellar Tendon-Bone and Hamstring Tendon Autografts for Anterior Cruciate Ligament Reconstruction: A Systematic Review and Meta-analysis. *Am J Sports Med.* 2022; 50: 3425-3439.
21. Sandon A, Werner S, Forssblad M. Factors associated with returning to football after anterior cruciate ligament reconstruction. *Knee Surgery, Sports Traumatology, Arthroscopy.* 2015; 23: 2514-2521.
22. Fältström A, Hägglund M, Kvist J. Factors associated with playing football after anterior cruciate ligament reconstruction in female football players. *Scand J Med Sci Sports.* 2016; 26: 1343-1352.
23. Burland JP, Toonstra JL, Howard JS. Psychosocial Barriers After Anterior Cruciate Ligament Reconstruction: A Clinical Review of Factors Influencing Postoperative Success. *Sports Health.* 2019; 11: 528-534.
24. Webster KE, Nagelli CV, Hewett TE, Feller JA. Factors Associated With Psychological Readiness to Return to Sport After Anterior Cruciate Ligament Reconstruction Surgery. *Am J Sports Med.* 2018; 46: 1545-1550.
25. Lisee CM, DiSanti JS, Chan M, Ling J, Erickson K, Shingles M, et al. Gender Differences in Psychological Responses to Recovery After Anterior Cruciate Ligament Reconstruction Before Return to Sport. *J Athl Train.* 2020; 55: 1098-1105.
26. Kunnen M, Dionigi RA, Litchfield C, Moreland A. Psychological barriers negotiated by athletes returning to soccer (football) after anterior cruciate ligament reconstructive surgery. *Ann Leis Res.* 2021; 26: 1-22.
27. Lentz TA, Paterno MV, Riboh JC. So you think you can return to sport? *Br J Sports Med.* 2018; 52: 1482-1483.
28. te Wierike SC, van der Sluis A, van den Akker-Scheek I, Elferink-Gemser MT, Visscher C. Psychosocial factors influencing the recovery of athletes with anterior cruciate ligament injury: a systematic review. *Scand J Med Sci Sports.* 2013; 23: 527-540.
29. Christino MA, Fantry AJ, Vopat BG. Psychological Aspects of Recovery Following Anterior Cruciate Ligament Reconstruction. *JAAOS - J Am Acad Orthop Surg.* 2015; 23: 501-509.
30. Fältström A, Kvist J, Gauffin H, Hägglund M. Female Soccer Players With Anterior Cruciate Ligament Reconstruction Have a Higher Risk of New Knee Injuries and Quit Soccer to a Higher Degree Than Knee-Healthy Controls. *Am J Sports Med.* 2018; 47: 31-40.
31. Sugimoto D, Myer GD, Barber Foss KD, Pepin MJ, Micheli LJ, Hewett TE. Critical components of neuromuscular training to reduce ACL injury risk in female athletes: meta-regression analysis. *Br J Sports Med.* 2016; 50: 1259-1266.
32. Jeon N, Choi NH, Hwangbo BH, Victoroff BN. An Increased Lateral Femoral Condyle Ratio in Addition to Increased Posterior Tibial Slope and Narrower Notch Index Is a Risk Factor for Female Anterior Cruciate Ligament Injury. *Arthroscopy.* 2022; 38: 1597-1604.
33. Crossley KM, Patterson BE, Culvenor AG, Bruder AM, Mosler AB, Mentiplay BF. Making football safer for women: a systematic review and meta-analysis of injury prevention programmes in 11 773 female football (soccer) players. *Br J Sports Med.* 2020; 54: 1089-1098.