

## Short Communication

# Evaluation of the Antischistosomal Effect of Poly I:C or Zymosan against *Schistosoma Mansoni* Infectious Disease

Mohamed Labib Salem<sup>1,2</sup>, Afrah Fathi Salama<sup>3</sup>, Wafaa Mohammed Ibrahim<sup>4</sup>, and Samah Kotb Nasr Eldeen<sup>3\*</sup>

<sup>1</sup>Zoology Department, Tanta University, Egypt

<sup>2</sup>Center of Excellence in Cancer Research, Tanta University New Teaching Hospital, Egypt

<sup>3</sup>Chemistry Department, Tanta University, Egypt

<sup>4</sup>Medical Biochemistry Department, Tanta University, Egypt

\*Corresponding author

Samah Kotb Nasr Eldeen, Chemistry Department, Tanta University, Egypt, Email: science\_samah@yahoo.com

Submitted: 23 January 2017

Accepted: 17 February 2017

Published: 20 February 2017

ISSN: 2378-931X

Copyright

© 2017 Nasr Eldeen et al.

OPEN ACCESS

## Keywords

- PolyI: C
- Zymosan
- SWAP antigen
- Adjuvant
- *S. mansoni*
- Vaccination
- Treatment

## Abstract

**Introduction:** In this study we focused on novel antischistosomal toll-like receptor (TLR) agonists which induced immune cells to produce a plethora of inflammatory cytokines against *Schistosoma mansoni* infection. We used in this study two important agonists TLR3L (polyinosinic/sidylic acid; poly I:C) and TLR2L zymosan. We divided this study into 2 experiments 1. The adjuvant effect of polyI:C or zymosan with soluble worm antigen (SWAP antigen) and we divided it into negative control group, positive control group, vaccinated group with SWAP antigen, vaccinated group with SWAP antigen plus poly I:C, vaccinated group with SWAP antigen plus zymosan 2. Antischistosomal effect of poly I:C or zymosan and we divided it into negative control group, two positive control groups, treated groups with poly I:C on day 25 or day 35 after infection, treated groups with zymosan on day 25 and day 35 after infection. Antischistosomal effect of poly I:C or zymosan was evaluated by measuring serum biochemical parameters as alanine aminotransferase, aspartate aminotransferase, alkaline phosphatase, gamma glutamyl transferase, albumin and total protein. We measured also oxidative stress parameters such as glutathione S. transferase, catalase, total antioxidant capacity, total thiol and malondialdehyde in liver homogenate, in addition to the histopathological examinations of all mice liver tissue with measuring granuloma sizes of all infected and treated groups and finally we measured the parasitological parameters as worm burden, liver and intestine eggs count.

**Results:** The results showed that polyI:C or zymosan with or without SWAP antigen improved liver enzymes, antioxidant parameters as well as it reduced the granuloma volume, worm burden, and ova count of all treated and vaccinated groups

**Conclusion:** PolyI:C or zymosan has adjuvant and antischistosomal effect against *S. mansoni* disease.

## INTRODUCTION

Schistosomiasis is a parasitic disease caused by platyhelminth worms of the genus *Schistosoma*. The disease affects 207 million people in the developing world, with approximately 800 million, mostly children, at risk of the infection [1-3]. No effective vaccine is available, despite the plethora of candidate vaccine antigens. Schistosomes are parasitic worms that are a prime example of a complex multicellular pathogen that flourishes in the human host despite the development of a pronounced immune response. The past decade has seen the use of a wide range of new approaches to determine the nature and function of the immune response to schistosomes. Infections with schistosomes are chronic and are characterized immunologically by an early T helper 1 response that switches to a T helper 2-dominated response after the onset of parasite egg production [3].

Toll-like receptors (TLRs) are pattern recognition receptors

that play a central role as sensors of infection and inducers of innate and adaptive immune responses [4] and are therefore considered as potential targets for vaccine adjuvants. They are the major class of signaling receptors which recognize pathogen associated molecular patterns (PAMPs) and signal the presence of an invading pathogen. They are members of a family of transmembrane proteins with an extracellular leucine-rich domain and a conserved cytoplasmic domain homologous to that of the interleukin-1 receptor (IL-1R), termed the Toll/IL-1R homology (TIR) domain. This structure allows TLRs to recognize PAMPs and activate, via the TIR domain, a series of downstream pathways that result in immune and inflammatory responses [5].

The specificity of TLR recognition has been established for several important PAMPs: bacterial lipoproteins and zymosan are recognized by TLR2 [6] double-stranded RNA by TLR 3 [5], lipopolysaccharide by TLR4 [5], flagellin by TLR5 [7], imiquimod by TLR7/8 and CpG motifs of bacterial DNA by TLR9 [8]. Ligation

of TLRs by TLR-ligands leads to induction of cytokine production and activation of immune system cells resulting in a better antigen presentation [9].

Given these immunostimulatory effects of TLR ligands, the aim of the present study was to investigate the adjuvant effect and antischistosomal activity of some toll like receptor ligands (Polyinosinic: polycytidylic acid TLRL3 and zymosan TLRL2). The criteria used for assessment the activity of the ligands included worm burden, tissue eggs count, oogram and histopathological findings. Since this endemic disease affects the liver, intestine, spleen and urinary bladder, it was of interest in the present study to assess the levels of total thiol, malondialdehyde (MDA), catalase, glutathione S. transferase and total antioxidant capacity in liver tissues homogenate, and serum liver functions (transaminases, alkaline phosphatase, gamma glutamyl transferase, total protein and albumin of all infected and treated groups.

## MATERIALS AND METHODS

### Chemicals

All chemicals used in the present study were high analytical grade products from Sigma (USA), InvivoGen (USA).

### Animals

Female Swiss albino mice, weighing 20-25 g were purchased from Theodor Bilharz Research Institute at Cairo, Egypt and maintained on balanced diet, free access to water and were let for about one week before experimentation to adapt the laboratory conditions.

### Ethics

Handling, anaesthetic and sacrifice procedures followed ethical guidelines approved by the Ethical Committee of the Federal Legislation, the National Institutes of Health Guidelines in the USA.

## EXPERIMENTAL DESIGN

The study included the use of poly I:C or zymosan as adjuvant with SWAP antigen. Mice were divided into 5 groups, each group included 10 mice, as following (1) negative group (2) positive control (3) mice were vaccinated with subcutaneous injection of 50 µg/200µL of SWAP antigen alone 2 times, 2 weeks intervals [10] (4) Mice were vaccinated with subcutaneous injection of 50 µg/200µL of SWAP antigen plus 25µg/300µL of poly I:C, then intraperitoneal (i.p) injection with 25µg/300µL of poly I:C 2 times, 2 weeks intervals [11] (5) Mice were vaccinated with subcutaneous injection of 50 µg/200µL of SWAP antigen plus 5 µg/200µL zymosan then intraperitoneal (i.p) injection with 5µg/200µL of zymosan 2 times, 2 weeks intervals [12] then all groups were euthanized on day 35 after *S. mansoni* infection. In the antischistosomal effect of poly I:C or zymosan, the study were divided into (1) negative control group (2) two positive control groups (3) treated groups with intraperitoneal (i.p) injection of 50 µg/300µL poly I:C on day 25 or day 35 after infection (4) treated groups with 10 µg/200µL zymosan (i.p) on day 25 and day 35 after infection, 2 times a week for 2 weeks and the mice were euthanized on day 39 or day 49.

## Blood specimens

At the end of each experiment mice were captured, blood samples were collected by eye bleeding. Collection of blood was from the orbital sinus of the mouse. The blood without anticoagulant was centrifuged at 3000r.p.m for 10 minutes in room temperature and serum was separated then kept at -20°C until assays. Serum was subjected to the following analysis: 1- Alanine aminotransferase (ALT) 2- Aspartate aminotransferase (AST) 3- Alkaline phosphatase (ALP) 4- Gamma glutamyl transferase (GGT) and 5- Albumin 6- Total protein.

## Tissue samples

Mice of all group were euthanized, dissected and after perfusion of their livers were rapidly excised. Accurately weighted tissue samples were divided into three parts, the first part for histopathology, the second part for egg count, the third part were homogenized in phosphate buffer PH (7.4) and were frozen at -20°C till usage for measurement of oxidative parameters. The enzyme activities were determined as followed 1- Glutathione S. transferase (GST) 2- Catalase and 3- Total antioxidant capacity (TAC) 4- Total thiol 5- Malondialdehyde (MDA).

## Preparation of tissue homogenates

Liver tissue was homogenized in 0.9N NaCl by a ratio 1:10 w/v for estimation of all enzymes under investigation.

## Parameter assays

Enzyme activities were evaluated using end point assay method. GST: the formation of the adduct, due to conjugation of GSH with 1-chloro-2, 4-dinitrobenzene (CDNB) the absorbance at 340 nm [13]. Catalase: 3ml buffered H<sub>2</sub>O<sub>2</sub> sample, mixed by the sample was read for 1min at 250 nm. MDA: was estimated by the method of [14] and was measured colourimetrically at 530 nm. Total Thiol: was performed by the method of [15] and was measured colourimetrically at 412 nm. TAC: At low pH, when a ferric tripyridyltriazine (FeIII-TPTZ) complex is reduced to the ferrous form (FeII), an intense blue colour with an absorption maximum at 593 nm develops and hence colour formation is the reducing ability of the sample according to [16]. AST and ALT were estimated through measuring oxaloacetate and pyruvate produced respectively [17]. ALP were measured as a liberated phenol in the presence of amino-4-antipyrine and sod-arsenate as a blocking agent and potassium ferricyanide as a color reagent. The developed color measured at 510 nm [18]. GGT: The substrate L-γ-glutamyl 4-nitroanilide, in the presence of glycylglycine is converted by γ GT in the sample to 4- nitroaniline which can be measured at 405nm. Total protein was estimated by the method of [19] using Coomassie Blue Dye in the present of bovine serum albumin. The developed color was measured at 595 nm. Albumin: in the presence of bromcresol green at a slightly acid PH, produces a colour change of the indicator from yellow - green to green - blue measured at 630.

## Measurement of granuloma volume

Lesions containing eggs in their centers were selected for measurement and the diameter of each liver granuloma was obtained by measuring two diameters of the lesion at right angles to each other using an ocular micrometer. The mean diameter of

all slide lesions from each mouse of each group was determined and the volume of each lesion was calculated, assuming a spherical shape [20], from its mean diameter using the following formula: Volume =  $R^3 \times \frac{22}{7} \times \frac{4}{3}$ .

## WORM BURDEN

Worms were recovered by liver perfusion as described by [21].

### Ova count

The number of ova/g tissue was counted by the method of [22], where

Number of ova in 1 g of liver = Number of ova in 5 ml KOH

Weight of liver in grams recorded before digestion in KOH

### Statistical analysis

Data in the present study are presented as mean  $\pm$  S.E. Statistical significance values were determined by one way analysis of variance (ANOVA) accompanied by (GraphPad in Stat Software).

## RESULTS

Significant increase in serum ALT, AST, ALP, GGT enzymes, albumin, total protein and MDA were observed after *S. mansoni* infection  $P \leq 0.05$ , whereas levels of catalase, GST, TAC and total thiol showed significant decrease  $P \leq 0.05$  compared to normal healthy mice. Vaccination with SWAP antigen, SWAP antigen plus polyI: C and SWAP antigen plus zymosan showed increased levels of catalase, GST, TAC and total thiol compared to infected animals  $P \leq 0.05$  whereas levels of ALT, AST, ALP, GGT enzymes, albumin and total protein showed significant decrease  $P \leq 0.05$

compared to infected animals (Table 1,2). Treatment with polyI: Corzymosan showed increased levels of catalase, GST, TAC and total thiol compared to infected animals  $P \leq 0.05$  whereas levels of ALT, AST, ALP, GGT enzymes, albumin and total protein showed significant decrease  $P \leq 0.05$  compared to infected animals (Table 3,4).

We noticed a significant decrease in worm burden in *S. mansoni*-infected mice vaccinated with SWAP antigen alone  $P \leq 0.01$  and SWAP antigen plus polyI: CP  $\leq 0.05$  compared to the *S. mansoni* -infected animals (Table 5). Treatment with polyI: C or zymosan showed decreased worm burden on both day 25 and day 35 compared to the *S. mansoni* -infected animals (Table 6). Vaccination with SWAP antigen  $P \leq 0.001$  or SWAP antigen plus polyI: C or SWAP antigen plus zymosan  $P \leq 0.01$  induced a significant decrease of egg count in 1 gram liver or intestine in comparison with the *S. mansoni* infected animals (Table 5). Treatment with polyI: C or zymosan showed decreased egg count on both day 25 and day 35 compared to the *S. mansoni* -infected animals (Table 6).

Histopathological examination of all groups showed **Negative control group**: normal liver cells (hepatocytes) radiating from a central vein with normal hepatic cords and normal portal tract (Figure 1). **Positive control group**: hepatic pathological significant changes induced that *S. mansoni*-infection resulted in formation of granulomas around the schistosome eggs. The granulomatous reaction was collagenized and formed of histocytes, lymphocytes, eosinophils and fibroblasts schistosomal pigments appeared as dark granules in the Kupffer cells (Figure 2). **Schistosomal infected animals after vaccination**: Mice vaccinated with SWAP antigen alone (Figure 3) induced insignificant reduction in granuloma volume in comparison

**Table 1:** Effect of the administration of poly I:Cor zymosan as adjuvant with SWAP antigen on serum parameters in **Group A:** Negative control **Group B:** Positive control **Group C:** SWAP antigen **Group D:** SWAP antigen plus PolyI:C **Group E:** SWAP antigen plus zymosan.

Parameters	Group A	Group B	Group C	Group D	Group E
ALT(U/L)	29.4 $\pm$ 4.71	43.25 $\pm$ 3.9*	48.3 $\pm$ 2.43*	44.05 $\pm$ 5.63*	24.6 $\pm$ 0.1 *
AST(U/L)	111.75 $\pm$ 15.8	144.8 $\pm$ 17.6*	195 $\pm$ 16.48 *	174.5 $\pm$ 27.3 *	67.8 $\pm$ 0.4*
ALP(U/L)	162.25 $\pm$ 4.71	177.63 $\pm$ 3.93	145.9 $\pm$ 2.43*	133.29 $\pm$ 5.63*	124 $\pm$ 0.1 *
GGT(U/L)	17.47 $\pm$ 0.81	59.78 $\pm$ 0.81*	15.45 $\pm$ 0.35*	16.25 $\pm$ 1.95*	26.69 $\pm$ 0.2*
Albumin(g/dl)	3.71 $\pm$ 0.12	3.14 $\pm$ 0.11	3.31 $\pm$ 0.09	3.45 $\pm$ 0.11	3.27 $\pm$ 0.02
Total protein (g/dl)	6.09 $\pm$ 0.16	5.41 $\pm$ 0.39	5.63 $\pm$ 0.23	5.95 $\pm$ 0.25	5.32 $\pm$ 0.52

Values represent mean  $\pm$  SE, n=10 for each group. The significance of difference was analyzed by one-way ANOVA using graphpadinstat Software  $P \leq 0.05$ .

**Table 2:** Effect of the administration of poly I:Cor zymosan as adjuvant with SWAP antigen on oxidative stress in **Group A:** Negative control **Group B:** Positive control **Group C:** SWAP antigen **Group D:** SWAP antigen plus PolyI:C **Group E:** SWAP antigen plus zymosan.

Parameters	Group A	Group B	Group C	Group D	Group E
Catalase (mmole/min/mg protein)	14.0 $\pm$ 1.0	7.49 $\pm$ 0.5 *	11.42 $\pm$ 0.6 *	11.5 $\pm$ 0.56*	12.1 $\pm$ 0.92*
GST( $\mu$ mole/min/mg)	38.7 $\pm$ 0.6	18.6 $\pm$ 1.7*	29.02 $\pm$ 2.4*	28.6 $\pm$ 1.6*	28.67 $\pm$ 1.23*
Total thiol( $\mu$ mole/g tissue)	55.3 $\pm$ 0.8	50.9 $\pm$ 0.8	42.8 $\pm$ 0.3*	64.7 $\pm$ 1.9*	49.2 $\pm$ 0.2 *
MDA(nmole/g tissue)	151.7 $\pm$ 12.9	226.2 $\pm$ 18.8*	164.5 $\pm$ 17.5*	167.4 $\pm$ 20.1*	165.4 $\pm$ 12.8*
Total antioxidant capacity ( $\mu$ mole/g tissue)	56.6 $\pm$ 6.3	29.2 $\pm$ 5.6*	61.0 $\pm$ 5.1*	60.1 $\pm$ 1.5*	69.9 $\pm$ 18.9*

Values represent mean  $\pm$  SE, n=10 for each group. The significance of difference was analyzed by one-way ANOVA using graphpadinstat Software  $P \leq 0.05$ .

**Table 3:** Effect of the administration of poly I:Cor zymosanas treatment on serum parameters in **Group A:** Negative control **Group B:** Positive control **Group C:** PolyI:C treated group (day 25), **Group D:** zymosan treated group (day 25) **Group E:** Non-infected animals **Group F:** Infected animals, **Group G:** Poly I:C treated group(day 35) **Group H:** zymosan treated group(day 35).

Parameters	Group A	Group B	Group C	Group D
	A. treatment on day 25 after infection			
ALT(U/L)	33.9±8.6	87.3± 1.1*	44.8± 9.3*	49.5±9.8*
AST(U/L)	68.6± 12.1	94± 2.6	65± 12.4	62± 3.7
ALP(U/L)	162.2±1.4	177.6 ±10.9	147.4± 5.4	142.7± 28.4
GGT(U/L)	31.4±0.1	59.7±9.2*	47.1± 9.3*	49.2± 0.8*
Albumin (g/dl)	3.71±0.12	2.1± 0.3	2.5± 0.1	2.0± 0.1
Total protein(g/dl)	6.09±0.16	3.4± 0.08*	4.4± 0.002*	4.6± 0.03*
B. treatment on day 35 after infection				
Parameters	Group E	Group F	Group G	Group H
ALT(U/L)	33.9±8.6	84.7± 1.9*	42.8± 9.9*	47.6± 9.2*
AST(U/L)	79.6± 12.1	95.3± 15.0	74.6± 15.1	70.0± 5.7
ALP(U/L)	162.2±1.4	182.2± 4.1	152.0± 2.6	139.9±9.7
GGT(U/L)	40.2± 1.0	59.7± 3.0*	50.6± 0.4*	58.9±0.8*
Albumin(g/dl)	3.71±0.12	2 ±0.09	3 ± 0.2	2.1± 0.5
Total protein(g/dl)	6.09±0.16	4.0± 0.4*	5.0 ± 0.4	4.4± 0.3

Values represent mean ± SE, n=10 for each group. The significance of difference was analyzed by one-way ANOVA using graphpadinstat Software P ≤ 0.05.

**Table 4:** Effect of the administration of poly I:Cor zymosanas treatment on oxidative stress (liver homogenate parameters)in **Group A:** Negative control **Group B:** Positive control **Group C:** Poly I:C treated group (day25), **Group D:** Zymosan treated group day 25 **Group E:** Non-infected animals, **Group F:** Infected animals, **Group G:** PolyI:C treated group day 35 **Group H:** Zymosan treated group(day35).

Parameters	Group A	Group B	Group C	Group D
	A. treatment on day 25 after infection			
Catalase(mmole/min/mg protein)	12.0±1.01	7.23± 0.64*	11.66± 0.48*	11.55±0.79*
GST(μmole/min/mg)	38.1±0.3	13.65±0.72*	28.46± 1.32*	25.35± 0.45*
Total thiol (μmole/g tissue)	41.2±5.5	27.05± 7.97*	50.44± 1.6*	46.23± 3.2*
MDA (nmole/g tissue)	116.9± 9.02	179.29±8.91*	116.86± 14.92*	115.19± 15.0*
Total antioxidant(μ mole/g tissue)	62.2±4.2	35.24± 2.67*	62.06± 0.93*	58.68± 1.56*
B. treatment on day 35 after infection				
Parameters	Group E	Group F	Group G	Group H
Catalase(mmole/min/mg protein)	12.00±1.01	5.59± 0.28*	12.11± 1.41*	11.85± 0.66*
GST(μmole/min/mg)	23.70± 1.46	5.76± 1.24*	23.37± 1.32*	23.30±0.67*
Total thiol (μmole/g tissue)	41.25±5.5	36.55± 5.02*	48.86± 5.52*	34.34± 5.02
MDA (nmole/g tissue)	92.59± 6.32	162.81± 5.48*	93.92± 6.48*	92.40±1.34*
Total antioxidant (μ mole/g tissue)	65.57±4.87	35.58±2.947*	64.82± 6.13*	64.96± 1.52*

Values represent mean ± SE, n=10 for each group. The significance of difference was analyzed by one-way ANOVA using graphpadinstat Software P ≤ 0.05.

with *S. mansoni* - infected mice. Mice vaccinated with SWAP antigen plus poly (I:C) did not induce any significant change in granuloma volume compared to mice vaccinated with SWAP antigen alone but induced significant reduction in granuloma volume as compared with *S. mansoni* - infected mice (Figure 4). Mice vaccinated with SWAP antigen plus Zymosan did not induce any significant change in granuloma volume compared to SWAP antigen alone or *S. mansoni* - infected mice (Figure 5). Treated mice with poly (I:C) on day 25 post infection induced a significant reduction in granuloma volume as compared with *S. mansoni*-

infected mice (Figure 6,7). The granulomatous reaction consists mainly of histocytes, lymphocytes and eosinophiles. *S. mansoni* worm trapped with venules of portal tract was noticed (Figure 8). Mice treated with Zymosan day 25 post infection induced a significant reduction in granuloma volume as compared with *S. mansoni* - infected animals. Inflammatory cells consist mainly of histocyte, lymphocytes and eosinophiles (Figure 9). Mice treated with Poly (I:C) day 35 post infection did not induce any significant change in granuloma volume as compared to schistosoma infected animals . Inflammatory cells consist mainly of histocyte,

**Table 5:** Direct effect of the administration of poly I:C or zymosan as adjuvant with SWAP antigen on worm burden and egg count.


Parameters	Negative control	Positive control	Vaccinated with SWAP antigen	Vaccinated with SWAP antigen with poly I:C	Vaccinated with SWAP antigen with zymosan
Worm burden	-	27.5 ± 1.52	**21.66 ± 1.69	*23.14 ± 1.44	24.75 ± 1.49
Egg count of 1g liver	-	8928.5±505.2	***5750±381.9	7071.4±658.72	6750±322.98*
Egg count of 1g intestine	-	9678.5±513.9	*7250±761.18	**6357.143±432.64	*6125±426.95

Values represent mean ± SE, n=10 for each group. The significance of difference was analyzed by one-way ANOVA and Tukey test using computer program. ANOVA was significant at  $P \leq 0.05$ . Tukey test was significant from corresponding value at \*  $P \leq 0.05$ , \*\*  $P \leq 0.01$  and \*\*\*  $P \leq 0.001$ .


**Table 6:** Direct effect of the administration of poly I:C or zymosan treatment on worm burden and ova count.

Parameters	Control	Infected alone	Poly(I:C)	Zymosan
	A. treatment in day 25 after infection			
Worm burden	-	19 ± 3.18	19.37± 2.08	19.8± 2.48
Ova count of 1g liver	-	11428.57± 1217.08	**7571.42± 702.27	**5785.71± 447.97
Ova count of 1g intestine	-	10214.29 ±1040.01	*6928.57± 560.91	*7583.33±472.87
B. treatment in day 35 after infection				
Worm burden	-	26.83± 2.78	*16± 2.48	*15.66± 2.05
Ova count of 1g liver	-	12187.5± 1747.55	*9166.66± 1588.8	*9166.66± 1184.6
Ova count of 1g intestine	-	7875 ±965.55	11000 ± 983.38	9300 ± 1383.87


Values represent mean ± SE, n=10 for each group. The significance of difference was analyzed by one-way ANOVA and Tukey test using computer program. ANOVA was significant at  $P \leq 0.05$ . Tukey test was significant from corresponding value at \*  $P \leq 0.05$ , \*\*  $P \leq 0.01$  and \*\*\*  $P \leq 0.001$ .




**Figure 1** Section in a liver of non-infected mice showing normal hepatocytes (A), central veins and portal tract (B). (H&E stain, X 100).




**Figure 3** Section in a liver of vaccinated mice with antigen alone before *S. mansoni* infection showing minimal granulomatous reaction (A) and dilated veins within portal tract (B) (H&E stain, X 100).




**Figure 2** Section in a liver of *S. mansoni*-infected mice (day 35) showing the characteristic bilharzial granuloma with heavy inflammatory cellular reaction surrounding the egg. (A), also dilated vein within portal tract was noticed (B) (H&E stain, X 200).




**Figure 4** Section in a liver of vaccinated mice with antigen with Poly(I:C) before *S. mansoni* infection showing minimal granulomatous reaction within portal tract (A) (H&E stain, X 100).



**Figure 5** Section in a liver of vaccinated mice with antigen with Zymosan before *S. mansoni* infection showing moderate granulomatous reaction (A) and dilated Portal tract (B) (H&E stain, X 100).



**Figure 6** Section in a liver of *S. mansoni*-infected mice ( day 25 post infection) showing the characteristic granuloma reaction with heavy inflammatory cellular reaction surrounding the calcified egg (A) composed mainly of histocytes, lymphocytes and few eosinophils (H&E stain, X 400).




**Figure 7** Section in a liver of *S. mansoni*-infected mice in (day 25 post infection) showing the characteristic granuloma with heavy inflammatory cellular reaction surrounding the egg (A) (H&E stain, X 100).

lymphocytes and eosinophiles with entrapped worms with veins of portal tract (Figure 12). Mice treated with Zymosan day 35 post infection induced no effect on granuloma volume as compared with infected mice (Figure 13). Hepatic pathological changes showed that *S. mansoni*-infection resulted in formation of granulomas around schistosoma eggs. Long standing granuloma (day 35) induced progressive fibrosis denoting healing process.


Schistosomal pigments appeared as dark granules in the Kupffer cells (Figure 6).

## DISCUSSION


The liver plays an important role in the vital activities of the body where its hepatocytes show differences in the localization and concentration of some enzyme systems [23]. Many of these enzymes served as marker enzymes for different cell organelles




**Figure 8** Section in a liver of *S. mansoni*-infected mice and treated with Poly(I:C) (day 25 post infection) showing multiple ova with minimal granulomatous reaction surrounding the eggs (A) (H&E stain, X 100).




**Figure 9** Section in a liver of *S. mansoni*-infected mice and treated with Zymosan (day 25 post infection) showing calcified schistosoma egg surrounded by minimal granulomatous reaction (A) (H&E stain, X 100).




**Figure 10** Section in a liver of *S. mansoni*-infected mice (day 35 post infection) showing the characteristic granulomatous reaction with heavy inflammatory cellular reaction surrounding the eggs (A) (H&E stain, X 100).



**Figure 11** Higher magnification of previous section of *S.mansoni*-infected mice (day 35 post infection) showing the characteristic granuloma reaction with heavy inflammatory cellular reaction surrounded the eggs (A) composed mainly of histiocytes, lymphocytes and few eosinophils (H&E stain, X 400).



**Figure 12** Section in a liver of *S.mansoni*-infected mice and treated with Poly(I:C) (day 35 post infection) showing calcified schistosoma egg surrounded by moderate granulomatous reaction (A) (H&E stain, X 100).



**Figure 13** Section in a liver of *S.mansoni*-infected mice and treated with Zymosan (day 35 post infection) showing three schistosoma eggs surrounded by healing granulomatous reaction (A) (H&E stain, X 100).

and any defect of them will be reflected to the enzyme activity itself. Hence, studying changes in these enzymatic activities could be helpful in evaluating the possible side effects of different treatments on different cell organelles after *S. mansoni* infection and the improvement occurring in such enzymes after treatment.

It is concerned to study transaminases enzyme activities

which showed a significant increase after infection. [24] attributed the increase of transaminase enzyme activities in mice serum to the decrease in hepatic cell population due to liver fibrosis or due to their release of the enzyme from the damaged livers into the circulation as a result of increased cell membrane permeability. The observed increase of AST was more manifested than that of ALT denoting that, although the latter is more specific for liver cells, yet it is less sensitive than AST in detecting liver cell damage [25]. Moreover, the presence of considerably more AST in human hepatic tissue indicated that the released ALT is too diluted in the extracellular compartment to cause significant increase in the ALT activity in *S. mansoni* patients. Therefore, variations in the release, destruction or excretion of the two enzymes or an unknown metabolism aberration are probably important contributory mechanisms [26].

In the present study, ALP enzyme activity in infected mice showed a significant increase. [24] and [25] observed an elevation in ALP activity in murine liver after *S. mansoni* infection. They attributed the increase in enzyme activity to the irritation of the liver cells by toxins or metabolic products of growing schistosomules, adult worms and eggs or due to increased loss of intracellular enzyme by diffusion through cell membranes which appears to act as a stimulus to the synthesis of more enzyme protein. Higher rates of formation would, in turn, increase the rate of diffusion and hence increase serum activity [27]. [28] mentioned a significant rise in liver ALP isoenzyme in patients having hepatosplenic schistosomiasis. [29] added that the elevation of ALP enzyme activity in *S. mansoni* infected human is of intestinal origin especially since *S. mansoni* is a disease which primarily affected the intestine, while this elevation is not of hepatic origin as it is observed in both patients of *S. mansoni* and hepatosplenomegaly disease. [30] showed by histochemical studies an increase in ALP activity in experimental infection with schistosomiasis and in the late stage of human lesions by liver biopsy. [31] attributed the increase in enzyme activity to the proliferation of bile ductules and bile canaliculi as a result of schistosomiasis by ultrastructural examination of the liver specimens. This result confirmed the observation of [32] who suggested that the response of the liver to any form of biliary tree obstruction is to synthesize more ALP. Concomitantly GGT enzyme was significantly increased by *S. mansoni* infection. This result confirmed the results reported of [33] who recorded that liver fibrosis caused a marked increase in the activities of serum GGT. GGT levels increase due to destruction of bile ducts. It has been indicated that GGT elevation was most likely secondary to hepatobiliary involvement by *Schistosoma mansoni* and may indicate chronicity [33]. As regard to albumin, Mice treated with Poly(I:C) and Zymosan on day 25 post infection showed significant increase in total protein level compared to infected mice. In the hepatic disease as a result of bilharzial infection, protein anabolism decreases while protein catabolism increases. [34] elucidated that malabsorption may be a contributing factor in decrease of protein synthesis through a defect in absorption of amino acids. Also, It has been explained that schistosomiasis showed a reduction in biologically active albumin mRNA results in decreased albumin synthesis and may be responsible in part for hypo albuminemia. In addition, increased globulin and collagen mRNA is associated with increased globulin and collagen synthesis during hepatic fibrosis [35].

It was reported that in parasitic disease, there is a complex and dynamic physiological relationship between the parasite and the antioxidant defense components of the host [36]. In schistosomiasis, granuloma macrophages isolated from hepatic, intestinal and pulmonary lesions were found to release significant amounts of superoxide radical and  $H_2O_2$  radicals [37]. The oxidative processes which occur upon contact with *S. mansoni* eggs trapped in the liver seem to proceed uncontrolled, since the enzymatic activities involved in superoxide radical and  $H_2O_2$  detoxification decrease drastically. Such events may be, at least in part, responsible for the pathology associated with schistosomiasis [38]. The infection with *S. mansoni* not only triggers the production of reactive oxygen species, but it also leads to the alteration of the antioxidant defense mechanism [39]. Concerning Catalase activity, the present work extended to investigate the activity of catalase that reveals a significant reduction in its activity. In agreement with this [38] showed that peroxide dismutation yield  $H_2O_2$  which is detoxified by catalase resulting in decrease in its activity for bilharzial infected mice. While catalase showed a significant increase in its activity in mice vaccinated and treated with Poly (I:C) or Zymosan. Toxic substances and free radicals elaborated by worms consume antioxidants and may affect the capacity of the liver to detoxify or naturalize the effect of the toxic endogenous and exogenous compounds. The data obtained in the present study showed that, lipid peroxides were elevated by *S. mansoni* infection. This coincides with [34] who found that lipid peroxides were elevated by *S. mansoni* infection throughout the different durations of infection, and [37] who found that the production of free radicals in the chain of biochemical reactions results in an increase in lipid peroxides, the present results confirmed these finding via elevation of lipid peroxidation products (MDA) that were considered as a marker of lipid membrane damage up on schistosomiasis infection [40]. Mice vaccinated or treated with Poly (I:C) or zymosan showed significantly decrease in MDA liver content when compared to infected mice. [41] reported that; there are two correlations between collagen deposition and production of MDA and 4-hydroxynonenal (HN6) by hepatic cells, it has been reported that TLR ligands (Poly(I:C) or Zymosan) exert their effects through over expression of fibrogenic cytokines and through the upregulation of procollagen-1 mRNA [42,43]. There is controversy concerning, glutathione-S-transferase, total antioxidant capacity. [44] pointed out that levels of glutathione-S-transferase and total antioxidant capacity were decreased in human and mice infected with schistosoma mansoni. Moreover, [45] found that the activity of glutathione-S-transferase decreased in *S. mansoni* infected mice. While it is [34] confirmed that glutathione-S-transferase was unaffected in livers of mice infected with *S. mansoni*. In the present study, Mice vaccinated or treated with Poly(I:C) or Zymosan showed significant increase in GST activity and total antioxidant capacity when compared to infected mice. Data obtained for GST activity and total antioxidant capacity is in agreement with [38] that reflects the positive role of these ligands on this enzyme activity [46]. Concomitantly total thiol, the results of total thiol content revealed a significant reduction resulting from oxidative stress due to schistosomiasis [38]. Mice vaccinated or treated Poly (I:C) or zymosan showed significant increase in total thiol when compared to infected mice in agreement with [47]. It has been reported that liver total

thiol was drastically depleted in bilharzial infected mice and this depletion may be due to the increased cytotoxicity of  $H_2O_2$  in endothelial cells as a result of inhibition of glutathione reductase [34].

The present results showed improvement of most biochemical parameters under investigation. This improvement due to vaccination was confirmed by the observed decrease in worm burden, ova count and its distribution pattern in the liver and intestine. This reduction may be due to the effect of the selected antigen with poly I: C and zymosan.

The histopathological picture of the livers revealed alteration of the normal hepatic lobe architecture characterized by heavy cellular infiltration in the liver of infected mice, fibrous exudation and infiltration of eosinophils and mononuclear cells. The most characteristic lesions were marked by the presence of important fibroblastic tissue and the atrophy of the lobules [48]. Granuloma formation results from a delayed hypersensitivity response generated by the host against antigens secreted by the eggs of parasite [49]. Parasitic granulomas in the liver of infected animals with a central area of eosinophils, lymphocytes and macrophages were surrounded by a capsule of connective tissues and fibroblasts [35].

In the present study, polyI: C or zymosan has a dual function to reduce the worm and ova count and to reduce the number and size of liver granulomas. This type of ligands has been suggested to play a significant role in cell activation and the modulation of granulomatous hypersensitivity.

In conclusion, In the present data TLR ligands decrease egg counts and granuloma volume that decrease the toxic substances and free radicals elaborated that improve antioxidant enzymes documented by positive correlation between antioxidant enzymes, egg count and granuloma volume. [50] reported that oxidative stress potentiates the dsRNA-induced IL-8 release through (NF- $\kappa$ B) activation, and this potentiation might be partly explained by the increased TLR3 expression.,  $H_2O_2$  plus Poly(I:C) increased the TLR3 expression compared with the basal condition.

## REFERENCES

1. Christensen NO, Gotsche G, Frandsen F. Parasitological technique for use in routine laboratory maintenance of schistosomes and for use in studies on the epidemiology of human and bovine schistosomiasis. Teaching Note, Danish Bilharziasis Laboratory. 1984.
2. Andrade ZA. Schistosomiasis and liver fibrosis. *Parasite Immunol.* 2009; 31: 656-663.
3. El Ridi RA, Tallima HA. Novel therapeutic and prevention approaches for schistosomiasis: review. *J Adv Res.* 2013; 4: 467-478.
4. Iwasaki A, Medzhitov R. Toll-like receptor control of the adaptive immune responses. *Nat Immunol.* 2004; 5: 987-995.
5. Salem ML, Diaz-Montero CM, El-Naggar SA, Chen Y, Moussa O, Cole DJ. The TLR3 agonist poly (I:C) targets CD8+ T cells and augments their antigen-specific responses upon their adoptive transfer into naïve recipient mice. *Vaccine.* 2009; 27: 549-557.
6. Ozinsky A, Smith KD, Hume D, Underhill DM. Cooperative induction of pro-inflammatory signaling by Toll-like receptors. *J Endotoxin Res.* 2000; 6: 393-396.



7. Smith KD, Andersen-Nissen E, Hayashi F, Strobe K, Bergman MA, Barrett SL, et al. Toll-like receptor 5 recognizes a conserved site on flagellin required for protofilament formation and bacterial motility. *Nat Immunol.* 2003; 4: 1247-1253.
8. Akira S, Hemmi H. Recognition of pathogen-associated molecular patterns by TLR family. *Immunol Lett.* 2003; 85: 85-95.
9. Bowie AG, Haga IR. The role of Toll-like receptors in the host response to viruses. *Mol Immunol.* 2005; 42: 859-867.
10. Maghraby SA, Shker KH, Zahran HG, El-Sherbiny M. In vivo, the immunological effects of *Fasciola gigantica* worms homogenate mixed with saponin on mice infected with *Schistosoma mansoni*. *J Med Sci.* 2007; 7: 724-731.
11. Taillardet M, Haffar G, Mondière P, Asensio MJ, Pléau-Pison T, Burdin N, et al. Toll-like receptor agonists allow generation of long-lasting antipneumococcal humoral immunity in response to a plain polysaccharidic vaccine. *J Infect Dis.* 2010; 202: 470-479.
12. Margaret M. Kelly, Kelly McNagny, David L. Williams, Nico van Rooijen, Lori Maxwell, Carol Gwozd, et al. The lung responds to zymosan in a unique manner independent of toll-like receptors, complement, and dectin-1. *Am J Respir Cell Mol Biol.* 2007; 38: 227-238.
13. Habig WH, Pabst MJ, Jakoby WB. Glutathione-S-transferase. The first enzymatic step in mercapturic acid formation. *J Biol Chem.* 249: 7130-7139.
14. Mesbah L, Soraya B, Narimane S, Jean PF. Protective effect of flavonoides against the toxicity of vinblastine cyclophosphamide and paracetamol by inhibition of lipid-peroxydation and increase of liver glutathione. *Haematol.* 2004; 7: 59-67.
15. Sedlak J, Lindsay RH. Estimation of total, protein-bound and non-protein sulfhydryl groups in tissue with Ellman's reagent. *Anal Biochem.* 1968; 25: 192-205.
16. Benzie FF, Iris I, Strain JJ. The Ferric Reducing Ability of Plasma (FRAP) as a Measure of "Antioxidant Power". The FRAP Assay. *Anal Biochem.* 1996; 239: 70-76.
17. Reitman S, Frankel S. A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *Am J Clin Pathol.* 1957; 28: 56-63.
18. Kind PR, King EJ. Estimation of plasma phosphatase by determination of hydrolysed phenol with amino-antipyrine. *J Clin Pathol.* 1954; 7: 322-326.
19. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976; 72: 248-254.
20. Cheever AW, Hoffmann KF, Wynn TA. Immunopathology of schistosomiasis mansoni in mice and men. *Immunol Today.* 2000; 21: 465-466.
21. Smithers SR, Terry RJ. The infection of laboratory hosts with cercariae of *Schistosoma mansoni* and the recovery of worms. *Parasitology.* 55: 695-700.
22. Cheever AW, Anderson LA. Rate of destruction of *Schistosoma mansoni* eggs in tissues of mice. *Am J Trop Med Hyg.* 1971; 20: 62-68.
23. Gumucio JJ, Chiamale J. Liver cell and liver function. In *The Liver Biology and Pathology*, Raven Press, NY. 1988; 931-947.
24. El-Aasar AA, El-Merzabani MM, Zakhary NI, Farag HI, Abdeen AM, Abd El-Salam I, et al. Biochemical and biophysical studies on schistosomal liver of mice. *Egypt J Bilh.* 1989; 11: 19-33.
25. Awadalla HN, Sherif AF, Shafei AZ, Khalil HA, Guirgis FK. Enzyme levels in homogenates of liver from mice infected with *Schistosoma mansoni* and from uninfected mice. *Int J Parasitol.* 1975; 5: 27-31.
26. Salah LA, Kheireldin AA, Mansour MM, Hussein F. Levels of some serum enzymes in patients with schistosomiasis. *J Trop Med Hyg.* 1976; 79: 270-274.
27. Wilkinson JH. The origin and fate of serum enzymes. In: *An Introduction to Diagnostic Enzymology*, Libbey & Co, London. 1962; 259-261.
28. Abdel-Rahman HM, El-Shanawani FM, Hassan MM, Salem M, El-Sahly AM. Alkaline phosphatase isoenzymes abnormalities in hepatic schistosomiasis. *Egypt J Bilh.* 1993; 15: 41-48.
29. Mansour MM, Farid Z, Bassily S, Salah LH, Watten RH. Serum enzyme tests in hepatosplenic schistosomiasis. *Trans R Soc Trop Med Hyg.* 1982; 76: 109-111.
30. Hunter S, El-Roody A, Helmi F, Mousa A. Histochemical studies on hepatic schistosomiasis before and after hycanthone therapy. *J Egypt Med Assoc.* 1973; 56: 54-68.
31. Mansy SS, Abdel-Hady AA, El-Badrawy N, Akl MM, Badawy A. Bile ductule proliferation in experimental hepatic schistosomiasis light and electron microscopic study. *Egypt J Bilh.* 1990; 12: 93-102.
32. Kaplan MM. Alkaline phosphatase. *Gastroenterology.* N Engl J Med. 1972; 286: 200-202.
33. Al Mofarreh MA, Al Akwaa AY, Al Mofleh IA. Gammaglutamyl transpeptidase activity in patients with schistosomiasis. *Saudi J Gastroenterol.* 2003; 9: 15-19.
34. Rizk M, Fayed TA, Badwy M, Sabaa-El-Regal N. Effect of different durations of *Schistosoma mansoni* infection on the level of some antioxidants in mice. *Med J of Islamic world Academy of Sciences.* 2006; 16: 25-34.
35. El-Banhawey MA, Ashry MA, El-Ansary AK, Aly SA. Effect of Curcuma longa or parziquantel on *Schistosoma mansoni* infected mice liver--histological and histochemical study. *Indian J Exp Biol.* 2007; 45: 877-889.
36. Connors VA, Deburon I, Granath WO. *Schistosoma mansoni*. Interlukine-1 increases phagocytosis and superoxide dismutase by hemocytes and decreases output of cercariae in schistosomus-susceptible *Biomphalaria glabrata*. *Exp Parasitol.* 1995; 80: 139-148.
37. Shaheen AA, Abd el-Fattah AA, Ebeid FA. Effects of praziquantel treatment on lipid peroxide levels and superoxide dismutase activity in tissue of healthy and *Schistosoma mansoni* infected mice. *Arzneimittelforschung.* 1994; 44: 94-96.
38. Gharib B, Abdallahi OM, Dessein H, De Reggi M. Development of eosinophil peroxidase activity and concomitant alteration of the antioxidant defenses in the liver of mice infected with *Schistosoma mansoni*. *J Hepatol.* 1999; 30: 594-602.
39. Pascal M, Abdallahi OM, Elwali NE, Mergani A, Qurashi MA, Magzoub M, et al. Hyaluronate levels and markers of oxidative stress in the serum of Sudanese subjects at risk of infection with *Schistosoma mansoni*. *Trans R Soc Trop Med Hyg.* 2000; 94: 66-70.
40. Bedossa P, Houghlum K, Trautwein C, Holstege A, Chojkier M. Stimulation of collagen 1(I) gene expression is associated with lipid peroxidation in hepatocellular injury. a link to tissue fibrosis. *Hepatology.* 1994; 19: 1262-1271.
41. Parola M, Leonarduzzi G, Robino G, Albano E, Poli G, Dianzani MU. On the role of lipid peroxidation in the pathogenesis of liver damage induced by long-standing cholestasis. *Free Radic Biol Med.* 1996; 20: 351-359.
42. Poli G, Parola M. Oxidative damage and fibrogenesis. *Free Radic Biol. Med.* 1997; 22: 287-305.

43. Casini A, Ceni E, Salzano R, Biondi P, Parola M, Galli A, et al. Neutrophil-derived superoxide anion induces lipid peroxidation and stimulates collagen synthesis in human hepatic stellate cells. role of nitric oxide. *Hepatology*. 25: 361-367.
44. Sheweita SA, Mostafa MH, Ebeid F, El-Sayed W. Changes in expression and activity of glutathione-S-transferase in different organs of *S. hematobium* infected hamsters. *J Biochem Mol Toxicol*. 2003; 17: 138-145.
45. Hassan FAM, Abed GH, Abdel-Samii MAZ, El-Din MOH. Antischistosomal activity of ginger aqueous extract against experimental schistosoma mansoni infection in mice. *J Biomark*. 2016; 2: 2.
46. Wang Y, Utzinger J, Xiao SH, Xue J, Nicholson JK, Tanner M, et al. System level metabolic effects of a *Schistosoma japonicum* infection in the Syrian hamster. *Mol Biochem Parasitol*. 2006; 146: 1-9.
47. Hamed MA, Hetta MH. Efficacy of *Citrus reticulata* and Mirazid in treatment of *Schistosoma mansoni*. *Mem Inst Oswaldo Cruz*. 2005; 100: 771-778.
48. Frontera E, Roepstorff A, Gázquez A, Reina D, Serrano FJ, Navarrete I. Immunohistochemical distribution of antigens in liver of infected and immunized pigs with *Ascaris suum*. *Vet Parasitol*. 2003; 111: 9-18.
49. Ali SA, Hamed MA. Effect of *Ailanthus altissima* and *Zizyphus spina-christi* on bilharzial infestation in mice: histological and histopathological studies. *J Appl Sci*. 2006; 6: 1437-1446.
50. Koarai A, Sugiura H, Yanagisawa S, Ichikawa T, Minakata Y, Matsunaga K, et al. Oxidative Stress Enhances Toll-Like Receptor 3 Response to Double-Stranded RNA in Airway Epithelial Cells. *Am J Respir Cell Mol Biol*. 2010; 42: 651-660.

#### Cite this article

Salem ML, Salama AF, Ibrahim WM, Nasr Eldeen SK (2017) Evaluation of the Antischistosomal Effect of Poly I:C or Zymosan against *Schistosoma Mansoni* Infectious Disease. *J Vet Med Res* 4(2): 1073.