

Research Article

Review on Control of Haemonchus Contortus in Sheep and Goat

Mola Selemon*

Jimma University College of Agriculture and Veterinary Medicine School of Veterinary Medicine, Ethiopia

*Corresponding author

Mola Selemon, Jimma University College of Agriculture and Veterinary Medicine School of Veterinary Medicine, Ethiopia, Email: molaselemon97@gmail.com

Submitted: 30 April 2018

Accepted: 25 June 2018

Published: 27 June 2018

ISSN: 2378-931X

Copyright

© 2018 Selemon

OPEN ACCESS

Keywords

- Sheep and goat
- Haemonchosis
- Haemonchus contortus
- Anemia
- Submandibular edema

Abstract

Small ruminant livestock, such as sheep and goats, are extremely susceptible to internal Parasites, especially gastrointestinal nematodes. *Haemonchus contortus* is a highly pathogenic, blood-feeding nematode of small ruminants and a significant cause of mortalities worldwide. Haemonchosis is a particularly significant threat in tropical, subtropical and warm temperate regions, where warm and moist conditions favors the free-living stages, but periodic outbreaks occur more widely during periods of transient environmental favorability. Barber's pole worm is singly the most important of all the gastrointestinal nematodes that affect the survival and productivity of shoat owned by rural poor farmers in the developing world. Barber's pole worm which is called *Haemonchus contortus* by its scientific name is highly pathogenic blood sucking parasite. *Haemonchus contortus* is highly pathogenic blood sucking parasite. Haemonchosis caused by *Haemonchus contortus* is a predominant, highly pathogenic and economically important disease of sheep and goats. *Contortus* is active mainly in warm, humid climates in the summer months. High levels of prevalence, intensity and abundance of these parasites were generally observed around the middle of the two rain seasons, with peaks occurring in May and September of the year. There are number of factors such as poor nutrition, concurrent disease, stress, overstocking, or pregnancy/lactation can cause a loss of immunity to parasites. Haemonchosis can be diagnosed based up on the characteristic clinical sign of anemia, Submandibular edema, weight loss, and ill thrift along with finding large numbers of eggs in the feces. It causes great loss in sheep and goat farm. To prevent this problem the owner or farmer and veterinarian should work effectively. They should be prevent pasture contamination on larval stage in early spring via timely and planned treatment strategies, rotate different doses of drugs on animal basis, avoid under dosing antihelminthic, utilize safe pasture treatment and move shames.

ABBREVIATION

ACSRPC: American Consortium for Small Ruminant Parasite Control; COWP: Copper Wire Particle; DNA: Deoxyribonucleic; FEC: Fecal Egg Count; GI: Gastrointestinal; GIN: Gastrointestinal Nematode; HU: Haramaya University; K Da: Kilo Dalton; L: Larva; MOA: Minister of Agriculture; NGO: Non- Governmental Organization; PCR: Polymerase Chain Reaction; PCV: Packed Cell Volume; PH: Power of Hydrogen; PPR: Per Parturient Rise; and Goat; Spp: Shoat Sheep Species; US: United States; US\$: United States Dollar.

INTRODUCTION

There are a lot of diseases of shoat, but none of them are found everywhere as or present as direct a threat to shoat as internal parasites [1]. Gastrointestinal nematodes (GIN) that infect sheep and goats include *Haemonchus contortus*, *Trichostrongylus*

colubriformis, *T. axei*, *Teladorsagia (Ostertagia) circumcincta*, *Cooperia* spp., *Oesophagostomum*, *Trichuris ovis*, *Strongloides papillosus*, and *Bunostomum*. These parasites can contribute to the overall problem of gastrointestinal parasitism of ruminants. Barber's pole worm which is called *Haemonchus contortus* by its scientific name is highly pathogenic blood sucking parasite which is the most prevalent and important in most regions of the US, especially in the southern [2].

Barber's pole worm is singly the most important of all the gastrointestinal nematodes that affect the survival and productivity of shoat owned by rural poor farmers in the developing world. This haematophagus parasite is infamous throughout the humid tropic subtropics, which is responsible for acute disease outbreaks with high levels of mortalities, particularly in young animals [2]. It is economically important parasites in India about 103 million US\$ was spent. *H.contorus* is prominent

parasites amongst the reports of anthelmintic resistance that has emerged in all countries of the world that produce small ruminants. This emergence of multiple anthelmintic resistances has provided a spur for research on alternative forms of control. Recent surveys in developing countries have identified many plants that are intended and have the potential to be used as anthelmintic against this parasite [1].

Haemonchosis can be diagnosed based up on the characteristic clinical signs of anemia, submandibular edema, weight loss, and ill thrift along with finding large numbers of eggs in the feces. Female *Haemonchus* produce approximately 5,000 eggs per day and caprine can be infected with thousands of these worms and these results in tens to hundreds of thousands of eggs being shed on to pasture by each animal each day. Because the life cycle is so short (< 3 weeks), this cycle of infection pasture contamination, reinfection and more pasture contamination can rapidly transform pastures in to very dangerous places for goats. This is especially true in a warm environment such as Georgia, because transmission of *H. contortus* occurs virtually year – round.

As other form of most parasitic diseases, haemonchosis is most severe in young animal during their first year on pasture. However, since immunity to GI nematodes in goats is slow to develop and is incomplete, even mature goats are at considerable risk. There are number of factors such as poor nutrition, concurrent disease, stress, overstocking, or pregnancy/lactation can cause a loss of immunity to parasites. It is well established that ewes and does lose much of their protective immunity to GIN around the time of kidding/ lambing causing the number of parasites infecting the does to increase. Subsequently, parasite egg production and contamination of the environment with infective larvae increases, creating a dangerous situation for the highly susceptible young kids. This phenomenon, known as the per parturient rise (PPR) is an extremely important part of the epidemiology of *Haemonchus* and must be considered when designing control programs [2].

OBJECTIVES

The objective of this review will be:

1. To provide the concise review on *haemonchus contortus*
2. To highlight control and prevention strategies on *haemonchus contortus*

LITERATURE REVIEW

Etiology

Phylum: Nematoda; Class: Secernentea; Order: Strongylida; Family: Trichostrongylidae; Genus: *Haemonchus*; Species: *Haemonchus contortus*; The majority of gastrointestinal strongyles of ruminants belong to the family Trichostrongylidae. The genus *Haemonchus* is in the sub – family of *Haemonchinae* and consists of four main species in domestic ruminants, namely, *H. contortus* (in ovine and caprine), *H. placei* and *H. similis* (in bovine) and *H. longistipes* (in dromedary). *Haemonchus contortus* is a cylindrical gastrointestinal nematode commonly known as the red stomach worm, the wire worm or the Barber pole worm [3]. Adult worms are found in the abomasum of goats

and sheep [4]. *Haemonchosis* caused by *Haemonchus contortus* is a predominant, highly pathogenic and economically important disease of sheep and goats [5]. These parasites are common blood feeders that caused anemia and reduced productivity and can lead to death in heavily infected animals [6].

According to [7] *H. contortus* is a highly pathogenic parasite of small ruminants, capable of causing acute disease and high mortality in all classes of stock affected. These parasites produce large numbers of eggs per parasite per day; this together with suitable climates of high temperature and rainfall ensures year – round, undisrupted development of larvae. *H. contortus* infection is so significantly high and of importance that it made the top ten list of the most common pathogenic nematodes of sheep and goats in the tropics and together with other nematodes and the second most common gastrointestinal infection that results in death of sheep in Malaysia [8]. Gastrointestinal nematode infection ranks highest on a global index, with *H. contortus* being of overwhelming importance [7].

Morphology

An adult *H. contortus* measures about 15 to 30 mm long, the male being shorter than the female. The morphological characteristics of *H. contortus* are a mouth capsule with a single dorsal lancet and two prominent cervical papillae in the esophageal area. The male parasite is characterized by its copulatory bursa formed of two large lateral lobes and a small asymmetrically positioned dorsal lobe [9]. Together with the two chitins spicules, which are inserted in the female genital opening during copulation, this part of the worm is important for identification [10]. Female's parasites have a reddish digestive tube filled with ingested blood, spirally surrounded by two white genital cords (ovaries) [11]. Eggs of strongyle type with a diameter ranging between 70-85 µm [12] (Figure 1).

One study which was conducted in HU indicates that important parasites of sheep and goats were intensified and those factors affecting the epidemiology of these parasites of small ruminants in eastern Ethiopia were elucidated. Those parasites are includes that *Haemonchus*, *Trichostrongylus*, *Nematodirus*, *Oesophagostomum*, *Fasciola* and *Paramphistomum* species were common helminthes of sheep and goats in this part of Ethiopia. According to the researcher from these GIT parasites *H. contortus* is the most prevalent, representing more than 60% of the total worm burdens recorded in tracer tests. Sheep and goats examined at abattoir mean *H. contortus* worm counts exceeded 4000 during the peak time. High levels of prevalence, intensity and abundance of these parasites were generally observed around the middle of the two rain seasons, with peaks occurring in May and September of the year. This confirmed that the weather conditions of the wet seasons were generally favorable for the development, survival and transmission of the free- living stages of nematodes [13].

Barbers pole worm is species most commonly encountered in small ruminants. The disease caused by this parasitic worm is haemonchosis, which is the most frequently observed gastrointestinal problem in tropical and sub-tropical regions of the world [14]. Like any other the parasitic diseases it is in a dynamic interaction with its environment and the host, the outcome of which depends on various intrinsic and extrinsic factors.




Figure 1 Morphology of male and female *H. contortus* Haemonchus contortus (Rudolphi, 1802) Cobb, 1898; A, anterior end of female; B, anterior end of male; C, vulva of female; D, bursa of male showing spicules.

Risk factors

Intrinsic Factors: *H. contortus* is a highly prolific, blood-feeding parasite with various strategies to escape adverse climatic conditions and immune reactions of the host. Its ability to produce large number of eggs during its lifetime provides *H. contortus* with an advantage over other parasites in that it can easily contaminate grazing areas and may survive in its small ruminant hosts through frequent and rapid re-infections. Variations in the degree of infectivity of different *H. contortus* isolates have been documented. A comparison in infectivity between *H. contortus* isolates from France with those from West Indies (Guadeloupe) in two breeds of sheep, namely the Black Belly and the INRA 401, has shown that the latter established between than the former in the Black Belly [15] suggesting that it is important to take in to account parasite genetic diversity in different agro0 ecological zones.

EXTRINSIC FACTORS

The environment

Climate and vegetation: Factors including temperature, rainfall, humidity and vegetation cover influence patterns of parasite development [16]. In most tropical and sub-tropical countries, environmental temperatures are permanently favorable for larval development. The ideal temperature for larval development of many nematode species in the microclimate of the pasture or vegetation is between 22°C and 26°C while the best humidity is close to 100% [17].

Desiccation from lack of rainfall kills eggs and larvae rapidly and is the most lethal of all climatic factors. A pasture larval assessment in Ghana and the use of tracer lambs in Kenya revealed that very few or no *H. contortus* infective larvae were available during dry periods while numbers of larvae were high in the rainy seasons and shortly after [18]. Below 5°C, movement

and metabolism of L3 is minimal favoring prolonged survival as these larvae are enclosed in a double sheath and thus unable to feed to continuously renew their energy [19].

Nutrition: There is substantial evidence for a beneficial role of a good plane of nutrition in the resistance or resilience of sheep to GI nematode infections [20]. Nutrition can influence the development and consequences of parasitism in three different ways: (1) it can increase the ability of the host to cope with the adverse consequences of parasitism (resilience), (2) it can improve the ability of the host to contain and eventually to overcome parasitism (resistance) by limiting the establishment, development and fecundity of the parasitism and/or (3) it can directly affect the parasite population through affecting the intake of certain antiphrastic drugs. Highly metabolizable protein diets have been shown to augment resistance of Ile de France and Santa Ines lambs against *H. contortus*. Well - fed animals can withstand the harmful effects of GI parasitism and remain reasonably productive and may require less anti -helminthic treatments when compared with undernourished animals [21]. Major problem in this respect is that haemonchosis is more prevalent in regions where the animal feed resources are very scarce and/or improperly managed and therefore insufficient to satisfy the demand throughout the year.

Nature of the host

Breed: Though it is still not clear how natural selection might shape patterns of immune responsiveness in terms of type and strength of response, different breeds of sheep express different susceptibility to gastro- intestinal parasitic infections. In this respect the Santa Ines [22], Barbados Black Belly and Texel [23]. Breeds of sheep are known to be more resistant to infection with *H. contortus* compared with Suffolk and Ile de France, INRA 401 and Suffolk breeds respectively [11]. This is evidenced by reductions in faecal egg count (FEC) and/or worm number, slower worm development and reduced fecundity. Genetic variations in the resistance to *H. contortus* with in sheep flocks have also been demonstrated and used in breeding schemes in Australia [24].

Age, sex and reproductive status: In addition to genetic factors, animals of different ages and sex respond differently to parasitic infections under similar management conditions [11]. Young animals are generally more susceptible to parasitic diseases than adults [25]. It is believed that lower resistance to disease in young ruminants is partly due to immunological hypo responsiveness, and is not simply a consequence of their not having been exposed sufficiently to pathogens to develop immunity. Innate immunity, often age-related, is also considered important in many cases. This may be due to physico-chemical differences in the gut environment in adult compared with young hosts [26]. On the other hand, previous exposure to *H. contortus* infection could result in enhanced resistance to subsequent infections [27]. Improved resistance to *H. contortus* was reported in second infections in Rhon and Merino land and Black Belly and INRA 401 lambs [28]. This may be due to the alteration of immunological and physicochemical mechanisms that while incapable of controlling the primary infection is nevertheless able to influence the challenge infections [15].

In a recent study, it was reported that male lambs excreted significantly higher number of faecal eggs, carried higher number of *H. contortus* worms and were more anemic than their female counterparts [28]. Earlier studies with *H. contortus* showed that castration enhanced the resistance of male lambs to the extent that FEC were lower than those of female lambs suggesting the existence of hormone related influences [29]. The phenomenon of the peri-parturient rise (PPR) in nematode egg output is also of great importance in the epidemiology of GIT nematodes of sheep. This is due to a temporary loss of acquired immunity to infection at around the time of parturition and during lactation and the PPR in FEC started 2 to 4 weeks before lambing and continued in to lactation in the post-parturition period [30].

Pathogenesis

Haemonchosis in sheep may be classified as hyper acute, acute, or chronic. In the hyper acute form, death may occur within one week of heavy infection without significant signs. This form of the disease is very rare and appears only in highly susceptible lambs. The acute form is characterized by severe anemia accompanied by edema bottle jaw [31]. Anemia is also characteristic of the chronic infection, often of low worm burdens and is accompanied by progressive weight loss [19].

The chronic form is the most commonly observed during natural infections. The lesions are associated to anemia resulting from blood loss. *H. contortus* is known to produce calcium and a clotting factor binding substance known as calreticulin enabling the parasite to feed easily on host blood and in so doing cause hemorrhagic lesions [32].

Life cycle

H. contortus is a blood sucking gastro-intestinal nematode parasite belonging to the family Trichostrongylidae. The typical life cycle being's when the adult female lays eggs that are passed out in the feces of the animal. Hatching of the eggs is controlled by temperature and humidity and also to an extent by the larva within the egg [33]. Eggs are passed out in the faeces of the mammalian host. The eggs hatch and develop in to the L1 and L2 juvenile stages in the faeces while feeding on bacteria. The L1 stage usually takes about 4-6 days to develop under temperatures in the range of 24°C - 29°C [34]. The L2 stage then develops in to the L3 stage, which is referred to as the filariiform infective larvae, by shedding its cuticle.

The L3 stage remains in its cuticle and crawls up the grass blades awaiting ingestion by a final host (the herbivore), which becomes infected post-ingestion. This larva settles in the abomasum where it sheds its cuticle and burrows in to the abomasum layer where it develops in to the adult stage L4 [35]. This larva in turn sheds its cuticle and develops in to the adult stage (L5). Male and female worms mate and live in the abomasum where they feed on the blood of the host. The life cycle reported here is extracted from [3].

An important phenomenon observed in the life cycle that has epidemiological implications is arrested larval development or hypobiosis. Hypobiosis is the temporary cessation of development of a nematode at a particular point in its parasitic development. Hypobiosis usually follows the onset of cold

autumn/winter conditions in the northern hemisphere or very dry conditions in the subtropics and tropics [36]. It is usually due to an unfavorable environmental stimulus, such as cold weather or dry conditions received by the free-living L3 prior to ingestion and usually coincides with onset of winter or very dries conditions. Arrested development can occur in the gut of sheep or on pasture and ensures survival of the nematode under adverse climatic conditions. Subsequent maturation of the larvae due to resumption of development known as the 'Spring rise', when favorable conditions return in the spring, leads to a rapid rise in infections levels or fecal egg counts in the sheep [37] (Figure 2).

Clinical signs

The clinical signs, mainly anemia, edema and loss of weight in association with reduced hematocrit values might be characteristic of haemonchosis in sheep [19]. At Post mortem, the abomasum appears edematous with petechial hemorrhages, occasional nodular developments and a rise in pH. Feces are well formed; diarrhea occurring only in infections complicated by the presence of such species as *Trichostrongylus* specie and *Cooperia* species. Labs are the most seriously affected members of the flock, but older sheep under stress also may have fatal anemia A heavy *Haemonchus* species infection (20,000 - 30,000 worms) can kill a sheep very quickly [38].


Diagnosis

Clinical signs: *Haemonchus contortus* infection is clinically diagnosed by anemia, dehydration, and sub mandibular internal fluid accumulation those results in the formation of a bottle neck, diarrhea and low packed-cell volume (PCV). Infection also results in retarded growth; weakness reduced reproductive performance, general illness and death [4].

Microscopic techniques: *Haemonchus* eggs were identified on the basis of morphology [39]. Species whose eggs exhibit similar morphological characters and therefore cannot be distinguished from each other can further be subjected to larval culture and identification of the third stage larvae (L3) [40]. Supplementary diagnosis is achieved through the use of microscopic techniques by the recovery of *H. contortus* eggs from stool samples. Because the eggs of many important genera are morphologically similar and therefore hard to identify to species level, a better way of delineate species is by larval culture and identification of 3rd stage larvae.

Demonstration of parasite eggs in faecal material can prove the presence of infection and is the most commonly used diagnostic method. Nevertheless, this method does not always reveal the presence of the parasite during low level of parasitic burden and pre-patent periods requiring repeated examinations. Host resistance to GIT helminthes also delays egg laying and a change in female worm size affects its fecundity. Hence, egg counts do not necessarily reflect the number of worms present. Other methods like measurement of parasite-specific antibodies can be used as supplementary diagnostic tools [41].

Serological techniques: Alternative diagnosis is based on serological techniques including ELISA [42]. According to Thekisoe et al., 2007 the biggest drawback of tests that rely on the detection of antibodies is their inability to distinguish between



past and present infections. Molecular techniques have proved more rapid and accurate than both microscopy and serology hence there has been recent movement towards the development of molecular assays for detection of helminthes infections.

PCR: PCR was first developed in the 1980's by Kary B. Mullis [43]. PCR for detecting trichostrongyle infections, including *H. contortus*, in ruminants was developed by [40]. The assay consisted of four genus-specific primer/probe sets enabling sensitive and specific amplification of target DNA from four trichostrongyle genera in the presence of multiple infections. Many other successful PCR-based assays have been developed over the past decade for identification and differentiation of strongly infections. However, PCR assays are more expensive as they include the use of expensive probes and cheaper methods of diagnosis are a necessity, justifying the need to develop a PCR assay for detection of *H. contortus* infections as a supplementary diagnostic tool.

Worm count and identification: According to Zajac et al., 2006 *Haemonchus* can be recovered from each of the infected abomasum in slaughterhouses is possible to count and morphologically identify [44].

TREATMENT

One of the challenges of *haemonchus contortus* in shoa is killing the animals in short period of time in acute case and other economic losses in chronic condition. At this time treatment is important Albendazole 5mg/kg, Fenbendazole 5mg/kg, oxfendazole 5mg/kg, levamisole 7.5mg/kg, Ivermectin 0.2 mg/kg, Moxidectin 0.2 mg/kg and closantel 10 mg/kg (Table 1).

PREVENTION AND CONTROL MEASURES OR STRATEGIES

The objective of most parasite control method is not to

completely avoid the parasites in livestock, but to keep the population under a threshold, above which it would otherwise inflict harmful effects on the host population. Any parasite control method aimed at minimizing a given parasitic population must consider the basic disease determinants briefly described above. The relative success or failure of any control strategies can be judged in terms of immediate and long term objectives, the ultimate goal being increase production, minimizing risks regarding drug resistance and consumer and environment associated problems. Generally, nematode control strategies can be directed against the parasite in the host and in the environment [45].

Methods to control *H. contortus* must attempt to break the life cycle of the worm; whether through anthelmintic, animal management, or pasture management [46]. Anthelmintic drugs that remove the parasite from the intestines are the most common method for managing *H. contortus*. Chemical anthelmintic are often used to combat haemonchosis, because they are cheap, simple and cost effective; but parasite resistance to them is growing [47,48]. Ivermectin as well as albendazole and fenbendazole (both benzimidazoles) have produced the highest levels of resistance, and resistance with levamisole and moxidectin is increasing [49,50].

Resistance to these drugs is high because each one uses a specific mechanistic pathway to kill *H. contortus*. For example, ivermectin binds to glutamate-gated channels in the worm's nervous system, opening them, paralyzing the worm and killing them through starvation [51].

Natural dewormers build the animal's resistance to the worm, increasing their ability to expel the adult nematodes before they are able to attach to the intestines. *H. contortus* has evolved to thrive in weak intestinal environments and therefore it is harder for the parasite to establish in healthy digestive systems [52]. For

Table 1: List of some recommended drugs against haemonchus in sheep.

Chemical group	Anthelmintic	Prescribed dose
IMIDAZOTHIAZOLES	Levamisole	7.5mg/kg
BENZIMIDAZOLE	Albendazole 5mg/kg	5mg/kg
	Fenbendazole 5mg/kg	5mg/kg
	Oxfendazole 5mg/kg	5mg/kg
MACROCYCLIC LACTONES(ivermectins)	Ivermectin	0.2mg/kg
	Moxidectin	0.2mg/kg
SALICYLANILIDES	closantel	10mg/kg
Source: (bowman, 1999)		

example, garlic (*Allium sativum*) is a known antibacterial that likely derives its success in treating for *H. contortus* by making the host's digestive tract healthier [54].

Treatment against *H. contortus* using chemical and natural anthelmintic and dewormers is one approach to managing haemonchosis. Another common practice is selective treatment. Some organizations are educating farmers to use the FAMACHA method to identify and treat only those with high haemonchosis levels using their eye color. Farmers can compare the color of their sheep's conjunctivae using a scientifically developed FAMACHA card, Redder eye colors.

Indicate healthier sheep, white colors indicate dangerous anemia levels [54-56]. Others are promoting the use of the McMaster's method, a veterinary practice for determining *H. contortus* egg counts in sheep feces using microscopes and grid-lined McMaster's slides [49].

Other preventative measures are being investigated to lower exposure to the nematode and to maintain low levels of infection. The application of multiple methods of *H. contortus* management is suggested for achieving the highest rates of resistance and resilience in flocks [47]. Two of these methods are pasture rotation and nutrient supplementation. Proper pasture rotation allows time for on-pasture larvae to die out before they can be reconsumed and for grasses to grow higher than the larvae can climb [57].

Alternate grazing of different host species and alternation of grazing and cropping are management techniques that can provide safe pasture and give economic advantage when combined with anthelmintic [58]. Studies in the wet tropical climates of several Pacific Island countries showed that peak larval concentrations of *H. contortus* and *Trichostrongylus* species occurred on pasture about one week after contamination, but fell to barely detectable levels within 9 weeks [59]. Based on this, a rotational grazing system was designed which has resulted in a significant reduction in faecal egg counts as well as the number of anthelmintic treatments needed per year.

However, in many parts of Africa, communal pastoral systems do not allow for regulated grazing as a means of lowering exposure to infective larvae on pasture. Growing human populations and livestock densities coupled with the frequent drought in some regions necessitate unregulated animal movement in search of green pasture and drinking water [60]. Exploitation of refugia through alternate grazing of cattle and sheep [58], or sheep and

goats [61] to reduce pasture levels of infective larvae or dilute populations of drug resistant strains of parasites strains of parasites could be of great value in any management program.

Studies show that protein and herb supplements improve the health of the digestive tract, lessening the effects of infection and increasing host resilience [62]. Most animals develop immunity against internal parasites, which keeps the parasites from reproducing but doesn't kill them. It is young animals that have not developed immunity and those animals whose immunities are compromised that are the most affected by *H. contortus* [52].

Once a sheep is infected, *H. contortus* impairs the animal's functions by reducing its voluntary food intake and the efficiency with which its food is used [63,64]. At both of these stages, before and during infection, heightened immunity is crucial to controlling parasitic stages within the host and thereby increasing resilience and resistance [47].

CONCLUSION AND RECOMMENDATIONS

Sheep is one of the most important animals that provide farmers with food and other products and by products. One of the most noteworthy antagonistic factors to sheep farming is parasitic diseases; especially those caused by nematodes. *H. contortus* is one of the most important endoparasites of sheep. Haemonchosis *contortus* cause great economic loss in shoats with causative agent of *H. contortus*. It takes 21 days after ingestion of its infective agent L3. Live in abomasum of shoats and cause irritation, each worm suck blood. Besides weight loss, anemic, bottle jaw, slow growth, poor wool, quality occurs. It can be identified by use of PCV, fecal egg count, worm count and identification and effective management of internal parasites cannot be accomplished by using only one management factor, it is a combination of factors that will produce the most effective defense against internal parasites. Anthelmintic should be used only to treat animals when necessary, and should be thought of a limited resource to be used sparingly. Proper pasture and animal management is a key component to managing internal parasites in sheep and goat operations plus vaccination, copper oxide, feeding condensed tannin containing forage and use parasite trapping fungi. Generally, the presences of risk factors like warm and humid, poor animal management and seasonal change leads to feed stress and other concurrent diseases that could favor the distribution of disease.

Therefore based on the above conclusion the following recommendations are forwarded:

- Animals should be kept in high plane of nutrition (management) especially sheep in poor body condition and young's in order to develop resistance against haemonchosis.
- Strategic deworming of the parasite should be focus on sheep at the beginning of the grazing period to prevent the contamination of the pasture and at end so as to prevent load of parasite in dry season.
- An appropriate control and prevention methods of haemonchosis should be designed like: rotational grazing, avoiding over flocking and zero grazing.

- Use clean feeders and continue to practice good hygiene in the pens and pasture.
- Use gravel or concrete in the feedlot area to break the worm life cycle and to prevent reinfestation.
- Try a mixed species grazing program using cattle and goats.

REFERENCES

1. SawlehaQadir 1, AK Dixit 1, Pooj Dixit. Use of medicinal plants to control *Haemonchus contortus* infection in small ruminants. *Veterinary World*. 2010; 3: 515-518.
2. Ray M. Kaplan. Responding to The Emergency of Multiple Drug Resistant *Haemonchus contortus*: Smart Drenching FAMACHA®. proceedings of the Kentucky Veterinary Medical Association Morehead Clinic Days Conference, Morehead, Kentucky. 2005; 4-5: Athens, Georgia,pp.
3. HALE M. Managing internal parasites in sheep goats. *Attr*. 2006; 1-8.
4. Inaam, El ad, Tayeb, Ibe1, Sa shadad, Hassan t. The effect of *Haemonchucontortus* infection treatment with ivermectin on drug - metabolizing enzymes. *Research Journal of Animal Veterinary Sciences*. 2007; 2: 123-128.
5. Mortensenm LL, Williamson LH, TerrillTH, Kircher R, Larsen M, Kaplan RM. Evaluation of prevalence clinical implications of anthelmintic resistance in gastro-intestinal nematodes of goats. *J Am Vet Med Assoc*. 2003; 223: 495-500.
6. Githigia SM, Thamsborg SM, Munyua WK, Maingi N. Impact of gastro intestinal helminthes on production in goats in Kenya. *Small Rum Res*. 2001; 42:21-29.
7. Maphosa V, Masika PJ, Bizimenyer A, ES, Eloff JN. *In-vitro* anthelmintic activity of crude aqueous extracts of *Aloe ferox*, *Leonotisleonurus* *Elephantorrhiza* elephantine against *Haemonchuscontortus*. *Tropical Animal Health Production*. 2010; 42: 301-307.
8. Cheah TS, Rajamanickam C. Epidemiology of gastro- intestinal nematodes of sheep in wet tropical conditions in Malaysia. *Tropical Animal Health Production*. 1997; 29: 165-173.
9. Morales G, pino LA. Ecoepidemiology of *Haemonchuscontortus* bahiensis, ecotype present in sheep of arid zones of Venezuela. *Memorias do Instituto Oswaldo Cruz*. 1987; 82: 359-369.
10. Bussieras J, Chermette R. *ParasitologyVeterinaries: Helminthology*. 1988.
11. Getachew T. Roles of eosinophil regulation of population *Haemonchus contortus*. *Sciences of Ecologies, Veterinary, Agronomy and Bioengineers Toulouse France*. 2007; 212: 50-80.
12. Kassai T. *Veterinary Helminthology*. Real Educational & Professional publishing Ltd. Butterworth - Heinemann. 1999.
13. Sissay MM. Epidemiology, Anthelmintic Resistance its Management, PhD thesis, Faculty of Veterinary Medicine Animal Science Department of Biomedical Sciences Veterinary Pubic Health Division of Parasitology Virology Uppsala, Sweden. 2007.
14. Achi YL, Zinsstag J, Yao K, Yeo N, Dorchie P, Jacquet P. Host specificity *Haemonchus Spp.* for domestic ruminants in the savannah in northern Ivory Coast. *Veterinary Parasitology*. 2003; 116: 151-158.
15. Aumont G, Gruner L, Hostache G. Comparison of the resistance to sympatric allopatric isolates of *Haemonchuscontortus* of Black belly sheep in Guadeloups. *Veterinary parasitology*. 2003; 116: 139-150.
16. Tembely S, Lahloukassi A, Rege JE, Sovani S, Diedhiou ML, Baker RL. The epidemiology of abomasal nematodes of sheep in Sweden, with particular reference to over-winter survival strategies. *Veterinary parasitology*. 1997; 122: 207-220.
17. Krecek RC, Groeneveld HT, Maritz JI. A preliminary study of the effect of microclimate on the third- stage larvae of *Haemonchuscontortus* *Haemonchusplacei* irrigated pasture. *International Journal for Parasitology*. 1992; 22: 747-752.
18. Wanyangu SW, Karimi S, Mugmbi JM, Bain RK. Availability of *Haemonchuscontortus* L3 larvae on pastur at K IBOKO: semi-arid warm agro-climatic zone in Kenya. *ActaTropica*. 1997; 68: 183-189.
19. Urquhart GM, Armour J, Duncan JL, Dunn AM, Jennings FW. *Veterinary parasitology*. 1996; 19-164. 2nd ed. Blackwell Science.
20. Bricarello PA, Amarante AF, Rocha RA, Cabral Filho SL, Huntley JF, Houdijk, JG, et al. Influence of dietary protein supply on resistance to experimental infections with *Haemonchuscontortus* in the Ile de France Santa Ines lambs. *Veterinary Parasitology*. 2005; 134: 99-109.
21. Knox MR, Torres-Acosta JF, Aguilar- Caballero AJ. Exploiting the effect of diatar supplementation of small ruminants on resilience end resistance against gastrointestinal nematodes. *Veterinary Parasitology*. 2006; 139: 385-393.
22. Amarante AF, Bricarello PA, Rocha RA, Gennari SM. Resistance of Santa Ines, Suffolk Ile de France sheep to naturally acquired gastrointestinal nematode infections. *Veterinary Parasitology*. 2004; 120: 91-106.
23. Good B, Hanrahan JP, Crowley BA, Mulchay G. Texel sheep are more resistant to natural nematode challenge than Suffolk sheep based on faecal egg count nematode burden. *Veterinary Parasitology*. 2006; 136: 317-327.
24. Woolaston RR, Baker RL. Prospects of breeding small ruminants for resistance to internal parasites. *International journal for parasitology*. 1996; 26: 845-855.
25. Colditz IG, Watson DL, Gray GD, Eady SJ. Some relationships between age, immune responsiveness resistance to parasites in ruminants. *International Journal for Parasitology*. 1996; 26: 869-877.
26. Mulcahy G, O'Neill S, Donnelly S, Dalton JP. Helminthes at mucosal barriers interaction with the immune system. *Advanced Drug Delivery Reviews*. 2004; 56: 853-868.
27. Gaulty M, Kraus M, Vervelde L, van Leeuwen MA, Erhardt G. Estimating genetic differences in natural resistance in Rhone Merino lsheep following experimental *Haemonchuscontortus* infection. *Veterinary Parasitology*. 2002; 106: 55-67.
28. Gaulty M, Schackert M, Hoffmann B, Erhardt G. Influence of se on the resistance of sheep lambs to an experimental *Haemonchuscontortus* infection. *Deutsche Tierarztliche Wochenschrift*. 2006; 113: 178-181.
29. Shaw KL, Nolan JV, Lynch JJ, coverdale OR, Gill HS. Effect of Weaning, supplementation gender on acquired immunity to *Haemonchuscontortus* in lambs. *International Journal for parasitology*. 1995; 25: 381-387.
30. Tembely S, Lahlou-Kassi A, Rege JEO, Mukasa-Mugerwa E, AnindoD, and Sovani, S. Breed season effects on the peri-specific loop-mediated isothermal amplification LAMP for diagnosis of trypanosomosis. *Acta Tropical*. 1998; 102: 182-189.
31. Taylor, MA, Hunt KR, Wilson CA. Quick JT. Clinical observation of diagnosis control of *haemonchuscontotus* infection in per parturient ewes. *vet. Roc*. 1990; 126: 555-556.
32. Suchura, Joshi P. Characterization of *Haemonchusconrtus* calreticulin suggests its role in feeding immune evasion by the parasite. *Biochemical Biophysical Acta*. 2005; 172: 293-303.
33. BOWMAN DDO. *Georgis' Parasitology for Veterinarians*. 7th Ed., 255 - 283.

34. Schnieder T, Heise M, Epe C. Genus-specific PCR for the differentiation of eggs or larvae from gastrointestinal nematodes of ruminants. *Parasitology Research*. 1999; 85: 895-898.
35. Harmon AF, Williams ZB, Zarlenga DS, Hildreth MB. Real-time PCR for quantifying *Haemonchus contortus* eggs potential limiting factors. *Parasitology Research*. 2007; 101: 71-76.
36. Gatongi PM, Prichard RK, Ranjan S, Gathuma JM, Munayau WK, Cheruiyot H, et al. Hypobiosis of *Haemonchus contortus* in natural infections of sheep goats in semiarid area of Kenya. *Veterinary Parasitology*. 1998; 77: 49-61.
37. Waller PJ, Bernes G, Rudby-Martin L, Ljungstrom BL, Rydzik A. (2004).
38. Shapiro, Pathology parasitology for veterinary technicians. Thomson Delmar learning. 2005; 170.
39. Valderrabano J, Delfa R, Uriate J. Effect of level of feed intake on the development of gastrointestinal parasitism in growing lambs. *Vet Parasitology*. 2002; 104: 327-338.
40. Von Samson-himmelstjerna G, Harder A, Schnieder T. The epidemiology of abomasal nematodes of sheep in Sweden, with particular reference to over-winter survival strategies. *Veterinary parasitology*. 2002; 70: 207-220.
41. Eysker, M. Ploeger HW, Values of present diagnostic methods for gastrointestinal nematode infections in ruminants. *Veterinary Parasitology*. 2000; 41, 211-225.
42. Prasad A, Nasir A, Singh N. Detection of anti-*Haemonchus contortus* antibodies in sheep by dot-ELISA with immunoaffinity purified fraction of ES antigen during prepatency. *Indian journal of Experimental Biology*. 2008; 46: 94-99.
43. Lattes S, Ferte H, Delaunay P, Depaquit J, Vassallo M, Vittier M, et al. *Trichostrongylus colubriformis* nematode infections in humans, France. *Emerging Infectious Diseases*. 2011; 17: 1301-1302.
44. Zajac M Anne, Garya. *Conboy Veterinary Clinical parasitology*. 7th ed. Black well publishing Company. U.K. 2006.
45. Larsen M. Prospects for controlling animal's parasite nematodes by predacious micro fungi. *Parasitology*. 2000; 120: 121-131.
46. Hepworth K, Neary M, Hutchens T. *Managing Internal Parasitism in Sheep Goats*. West Lafayette, IN: Purdue University Cooperative Extension Service. 2006; 1-10.
47. Stear MJ, Doligalska M, Donskow-Schmelter K. Alternatives to Anthelmintics for the control of nematodes in livestock. *Parasitology*. 2007; 134: 139-151.
48. Yadav CL, Kumar R, Uppal RP, Verma SP. Multiple anthelmintic resistances in *Haemonchus contortus* on a sheep farm in India. *Vet Parasitology*. 1995; 60: 355-360.
49. Burke JM, Miller JE, Brauer DK. The effectiveness of copper oxide wire particles as an anthelmintic in pregnant ewes safety to offspring. *Veterinary Parasitology*. 2005; 131: 291-297.
50. Shoenian S, Slowing dewormer resistance. *Sheep and Goat.com*. Maryl Small Ruminant Page. 2013 <http://www.sheepandgoat.com/articles/slowdrugresist.html>.
51. Nolan. (2004) ivermectin. University of Pennsylvania. <http://cal.vet.upenn.edu/projects/dxendopar/drug%20pages/ivermectin.htm>
52. Wells A. Sustainable Management of Internal Parasites in Ruminants. NODPA News.
53. Quarterly Publication. Northeast Organic Dairy Producers Alliance, 2005; 20-23.
54. Noon J. A Controlled Experiment to Measure the Effectiveness on Lambs of Wormers. That Conform to the New Organic Standards. Sustainable Agriculture Research Education (SARE). Northeast SARE 2003 Farmer/Grower Grant Report.
55. Hart SP, Dawson LJ. Using FAMACHA alternative dewormers to manage gastrointestinal nematodes in a dairy goat herd. *Journal of Animal Science* 88 (E-Supplement 2), 2010; 580.
56. Van Wyk JA, Bath GF. The FAMACHA system for managing haemonchosis in sheep goats by clinically identifying individual animals for treatment. *Veterinary Research*. 2002; 33: 509-529.
57. Morgan, J. A Friendly Encouraging Challenge to the Agricultural Extension Community: A low cost tool that can greatly influence management of internal parasites in Small ruminants. *Sheep Farm Life*. 2005; 50: 34-35.
58. Machen R, Craddock, F, Craig T, Fuchs T. (1998). *A Haemonchus contortus Management Plan for Sheep Goats in Texas*. Pamphlet L-5095. College Station, T.X.: AgriLife Communications, Texas A&M System.
59. Barger Ian. Control by management. *Veterinary Parasitology*. 1997; 72: 493-506.
60. Bang KS, Familton AS, Sykes AR. Effect of ostertagiasis on copper status in sheep: a study involving use of copper oxide wire particles. *Res Vet Sci*. 1990; 49: 306-314.
61. Abule E, Snyman HA, Smit GN. Comparisons of pastoralists' perceptions about rangeland resource utilization in the Middle Awash Valley of Ethiopia. *Journal of Environmental Management*. 2005; 75: 21-35.
62. Sissay MM, Asefa A, Uggla A, Waller PJ. Anthelmintic resistance of nematode parasite of small ruminants in eastern Ethiopia: Exploitation of refugia to restore anthelmintic efficacy. *Veterinary Parasitology*. 2006; 135: 337-346.
63. Williams AR. Immune-mediated pathology of nematode infection in sheep – is immunity beneficial to the animal? *Parasitology*. 2011; 138: 547-556.
64. Van Houtert MF, Sykes AR. Implications of Nutrition for the Ability of Ruminants to Withstand Gastrointestinal Nematode Infections. *International Journal of Parasitology*. 1996; 26: 1151-1167.
65. Parkins JJ, Holmes PH. Effects of Gastrointestinal Helminth Parasites on Ruminant Nutrition. *Nutrition Research Reviews*. 1989; 2: 227-246.

Cite this article

Selemo M (2018) Review on Control of *Haemonchus Contortus* in Sheep and Goat *J Vet Med Res* 5(5): 1139.